

Chemistry

Manuscript Number: FOODCHEM-D-20-01584R2

Title: A Systematic Review of Phytochemicals in Oat and Buckwheat

Article Type: VSI: 4-ISPMF

Keywords: Oats; buckwheat; phytochemicals; bioactives; systematic review

Corresponding Author: Dr. Marija Glisic, PhD

Corresponding Author's Institution: Institute of Social and Preventive Medicine (ISPM)

First Author: Peter Francis Raguindin, MD

Order of Authors: Peter Francis Raguindin, MD; Oche Adam Itodo, MSc; Jivko Stoyanov, PhD; Gordana M Dejanovic; Magda Gamba; Eralda Asllanaj; Beatrice Minder; Weston Bussler; Brandon Metzger; Taulant Muka; Marija Glisic, PhD; Hua Kern

**Abstract:** Consumption of oat and buckwheat have been associated with various health benefits that may be attributed to their nutritional composition. We performed a systematic review to evaluate the profile and quantity of bioactive compounds present in oat and buckwheat. Among 154 studies included in final analysis, 113 and 178 bioactive compounds were reported in oat and buckwheat, respectively. Total phytosterols, tocopherols, flavonoids and rutin content were generally higher in buckwheat,  $\beta$ -glucans were significantly higher in oat, while avenanthramides and saponins were characteristically present in oat. The majority of studies included in current review were published before 2010s. The heterogeneous methodological procedures used across the studies precluded our possibility to meta-analyse the evidence and raises the need for harmonization of separation and extraction methods in future studies. Our findings should further stimulate the exploration of metabolites related to identified phytochemicals and their roles in human health

## Highlights

- Oat and buckwheat are good sources of micro- and macronutrients and bioactive phytochemicals, with 113 and 178 phytochemicals reported respectively.
- Compared to oat, total phytosterols, tocopherols, flavonoids and rutin content were generally higher in buckwheat.
- $\beta$ -glucans were significantly higher in oat while avenanthramides and saponins were characteristically found in oat.
- Proper food transformation might be a prerequisite to preserve nutritional content of oat and buckwheat.
- Bioactivity and bioavailability of oat and buckwheat's phytochemicals and their health effects need further investigation.

# A Systematic Review of Phytochemicals in Oat and Buckwheat

Peter Francis Raguindin<sup>1, 2\*</sup>, Oche Adam Itodo<sup>1,2\*</sup>, Jivko Stoyanov<sup>2</sup>, Gordana M. Dejanovic<sup>3</sup>, Magda Gamba<sup>1</sup>, Eralda Asllanaj<sup>4</sup>, Beatrice Minder<sup>5</sup>, Weston Bussler<sup>6</sup>, Brandon Metzger<sup>6</sup>, Taulant Muka<sup>1</sup>, Marija Glisic<sup>1,2¥</sup>, Hua Kern<sup>6¥</sup>

¥ and \* denote equal contributions

<sup>1</sup>Institute of Social and Preventive Medicine (ISPM), University of Bern, Bern, Switzerland

<sup>2</sup>Swiss Paraplegic Research, Nottwil, Switzerland

<sup>3</sup>Department of Ophthalmology, University of Novi Sad, Faculty of Medicine, Hajduk Veljkova 1-3, 21000 Novi Sad, Serbia

<sup>4</sup>Department of Epidemiology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, the Netherlands

<sup>6</sup>Standard Process Nutrition Innovation Center, Kannapolis, NC 28018, USA

<sup>5</sup>Public Health & Primary Care Library, University Library of Bern, University of Bern, Bern, Switzerland

<sup>6</sup>Standard Process Nutrition Innovation Center, Kannapolis, NC 28018, USA

Word count: Abstract: 145; Full text without abstract, references and figures: 3,554

Number of figures: 2

Number of tables: 2

Corresponding Author:

Marija Glisic, MD, PhD

Institute of Social and Preventive Medicine (ISPM), University of Bern, Bern, Switzerland

Email: marija.glisic@ispm.unibe.ch; Phone: +49 421 218 56845

## ABSTRACT

Consumption of oat and buckwheat have been associated with various health benefits that may be attributed to their nutritional composition. We performed a systematic review to evaluate the profile and quantity of bioactive compounds present in oat and buckwheat. Among 154 studies included in final analysis, 113 and 178 bioactive compounds were reported in oat and buckwheat, respectively. Total phytosterols, tocopherols, flavonoids and rutin content were generally higher in buckwheat,  $\beta$ -glucans were significantly higher in oat, while avenanthramides and saponins were characteristically present in oat. The majority of studies included in current review were published before 2010s. The heterogeneous methodological procedures used across the studies precluded our possibility to meta-analyse the evidence and raises the need for harmonization of separation and extraction methods in future studies. Our findings should further stimulate the exploration of metabolites related to identified phytochemicals and their roles in human health.

Key words: oats; buckwheat; phytochemicals; bioactives; systematic review

## 1. INTRODUCTION

Cereal grains are the major source of energy and an important constituent of diet for millions of people worldwide (Gangopadhyay, *et al.*, 2015); however, the majority of acreage dedicated to growing cereal crops throughout the world produce wheat, rice, and maize (Food and Agriculture Organization, 2003). This translates into consumption patterns that favor these crops over the other cereals, such as oat (*Avena sativa* L) and buckwheat (*Fagopyrum esculentum* Monch), that are less often cultivated worldwide.

Oats and buckwheat grains are unique due to their blend of macro, micro, and phytonutrients and their consumption have been associated with various health benefits including reduced risk of cancer, obesity, diabetes and cardiovascular diseases (CVD) (Borneo & Leon, 2012; L Li, *et al.*, 2018; Rasane, *et al.*, 2015). The observed health benefits have been attributed to their high content of dietary fibers such as  $\beta$ -glucan, functional protein, essential fatty acids, vitamins, and antioxidant phytochemicals, including high phenolic compounds (Flander L, 2007; Gangopadhyay, *et al.*, 2015). Due to its well-balanced nutritional profile, oat is one of the most important cereal crops in the developed world. Compared with other cereals, oats have higher concentrations of certain nutrients and phytochemicals and can tolerate harsher growing conditions, such as wet climate and acidic soil, making them more resilient than other crops (Grundy, *et al.*, 2018). Similarly, buckwheat, besides high starch content, is also rich in protein with a well-balanced amino acid profile, dietary fiber, lipids, and minerals, along with other components such as phenolic compounds and sterols (Ahmed, *et al.*, 2014; Krkoskova & Mrazova, 2005). Due to low gluten content, both oat and buckwheat are suggested to be suitable substitutes for wheat, barley, and rye in gluten-sensitive patients (Arendt & Zannini, 2013).

Recent studies suggested important differences in major phytochemical content between oat and buckwheat (Dar, *et al.*, 2018; Keriene, *et al.*, 2015; Yu, *et al.*, 2018). Previous attempts to review the

literature on the phytochemical content of oat (Boz, 2015; Gangopadhyay, *et al.*, 2015; Sang & Chu, 2017) and buckwheat (Ahmed, *et al.*, 2014; Zhan-Lu, *et al.*, 2012), were not systematic, did not report the quantities of compounds identified, and did not directly compare phytonutrients between the two cereals. Advancing the understanding of the two grains together can facilitate comparisons between oats and buckwheat, help support the utilization of the two grains for consumer products, and justify their combination in future studies. Thus, this systematic review of the literature aims to evaluate and compare the profiles and quantities of phytochemicals commonly found in oat and buckwheat, and to explore the potential of incorporating these two plants into human diets for optimal health and provide perspectives for further investigation. We hope the present systematic review will help present the two plants as a good option for growers to consider for their reported phytochemical contents and associated health benefits, and the benefit of a more diverse set of crop cultivation from our current mono-crop centric society.

## 2. METHODS

### 2.1 Literature search

This review was conducted following a recently published guideline on how to perform a systematic review (Muka, *et al.*, 2020), following the PRISMA guidelines (Moher, *et al.*, 2009) (Supplemental Table 1) and in accordance with the review protocol. Four bibliographic databases (PubMed, Embase, Web-of-Science, and Cochrane Central Register of Controlled Trials) were used to identify published studies until June 1, 2020 (date last searched) that examined the nutrient and bioactive composition of oat and buckwheat. The search terms we used were related to nutrient and bioactive compounds (e.g., nutrients, metabolism, phytochemical, carbohydrate, fatty acids) and the plants (oat and buckwheat) (Supplemental Table 2). We did not apply any restrictions on language and date. However, we excluded conference

abstracts, letters to the editor and editorials. To retrieve additional publications, we checked the reference lists of studies included in the current review.

## *2.2 Study selection criteria*

Studies were included if they met the following inclusion criteria: (i) used samples of oat and/or buckwheat; and (ii) evaluated nutrient and bioactive compounds. Two reviewers independently assessed the titles and abstracts according to the selection criteria. For each potentially eligible study, two reviewers assessed the full-text articles. In cases of disagreement, the decision was made by consensus, or, if necessary, a third reviewer was consulted.

## *2.3 Data extraction*

Two reviewers extracted data independently using a predesigned form, including first author and publication year, plant's source, compound name, and concentration, as reported in the articles.

## **3. RESULTS**

After searching the following electronic databases PubMed, Embase, Web of-Science, and Cochrane, 3,250 potentially relevant citations were identified, and after removing 817 duplicates, 2,433 abstracts and titles were evaluated according to inclusion and exclusion criteria (Figure 1). Full texts and reference lists of 201 studies were evaluated, among which 154 studies were eligible to be included in the current review. In the 154 included studies (oat n=72 and buckwheat n=83), 113 phytochemicals and metabolites were reported in oat and 178 in buckwheat, which we discuss in detail in the later sections. The list of all phytochemicals reported in oat and buckwheat and the references can be found in Table 1&2, while detailed information on phytochemical concentrations can be found in Supplemental Table 2 & 3. An illustrative summary of the most important findings is presented in Figure 2.

### 3.1 Phenolic Compounds

Phenolic compounds consist of one or more aromatic rings, bearing one or more hydroxyl groups, and are generally categorized as flavonoids, phenolic acids, stilbenes, polyphenolic amides (e.g., avenanthramides), coumarins, and tannins (Ahmed, *et al.*, 2014). Details on phenolic compounds identified in oat and buckwheat are summarized in Table 1 & 2 and Supplemental table 3.

#### 3.1.1. Flavonoids

Flavonoids (flavonols, flavones, flavanols, flavonones, flavans, and anthocyanins) are phenolic compounds, which comprise more than 4000 polyphenolic complexes (Tanwar & Modgil, 2012). In total, 16 and 64 flavonoids were reported in oat and buckwheat, respectively, and those are listed in Table 1 & 2 and Supplemental Table 2. Total flavonoid content ranged from 6 mg of rutin equivalents (RE)/100g in groat to 3,149 mg RE /100g in unfermented buckwheat sprouts (Chen, *et al.*, 2018; XD Guo, *et al.*, 2011; Peng, *et al.*, 2017; Sedej, *et al.*, 2012). Tartary buckwheat had a higher content of total flavonoids in comparison with common buckwheat, with an indication that this difference decreases during germination (Fabjan, *et al.*, 2003; Jiang, *et al.*, 2015; Tien, *et al.*, 2018). Rutin was a predominant flavonoid compound detected in buckwheat. Although its content was high in all morphological parts of buckwheat, its highest content was observed in buckwheat leaves (maximum 3,417 mg/100g) and bran (maximum 5,186mg/100g)(Habtemariam, 2019). Rutin levels were also up to 5-fold higher in tartary in comparison to common buckwheat (Y Liu, *et al.*, 2019). Lower concentration of rutin was also reported in husked oat fractions (0-3.2 mg/100g) and oat seeds (0.22-0.47 mg/100g) (Kerene, *et al.*, 2015; Tong, *et al.*, 2014). Other flavonoids such as vitexin, isovitexin, orientin, and isoorientin were generally more abundant in common buckwheat. Quercetin was the predominant flavonoid compound in oat with husked oat samples having the highest concentration (up to 8.9 mg/100g) (Kerene, *et al.*, 2015). In common buckwheat, quercetin concentrations ranged from 0.07 mg/100g in the hull to 33 mg/100g of buckwheat



sprouts with tartary buckwheat seeds reaching the levels as high as 291 mg/100g (XD Guo, *et al.*, 2011). In general, a wide range of variations of flavonoids among buckwheat and oat cultivars was linked with the variety, environmental factors, growing conditions, time of harvest, processing techniques and anatomical part of the plant used (Fabjan, *et al.*, 2003; XD Guo, *et al.*, 2011; B. D. Oomah & G. Mazza, 1996; Tong, *et al.*, 2014). Fagopyrin, flavonoid potentially harmful to human health, was mostly reported in buckwheat sprouts and ranged from 0.0025% to 0.041% and from 0.10% to 0.12 % in common and tartary buckwheat sprouts respectively; while in another study its concentration ranged from 19 to 32 mg/100g of common and tartary buckwheat leaves and flowers respectively (Habtemariam, 2019; Kreft, *et al.*, 2013). Kreft *et al.*, however, suggest that the consumption of approximately 40 sprouts (0.14 g of dry mass sprouts per kg of body mass per day) may not cause phototoxic effects. The intake of 10 g of dry mass (or approximately 30 g of fresh mass) of buckwheat sprouts may, on the other hand, cause health problems (such as dermatitis and hair loss). Yet, the authors warn that the toxic dose may vary depending on exposure to sunlight, body mass, and age (Kreft, *et al.*, 2013).

### 3.1.2. Polyphenolics

Total phenolic content in oat varied from 35.1 to 576 mg of mg/100g with oat bran concentrate having the highest content. In buckwheat, total phenolic content was expressed in gallic acid (GAE) or rutin (RE) equivalents reaching 670 GAE mg/100g in buckwheat seeds and sprouts, and 1,410 mg RE mg/100g in common buckwheat husks. In general, oat bran, hulls, and groats had the highest total phenolic content. Only two studies directly compared the total phenolic content in oat and buckwheat. Buckwheat husks and grain with husks had 2 to 5-folds more phenolic compounds in comparison to oats samples, as expressed in RE (Supplemental Table 2). In line with this, in another study comparing the GAE in both plants, the buckwheat had higher total phenolic content (330.3-3.951.4 mg GAE/100g) in comparison to oat (113.8 mg GAE/100g) (Holasova, *et al.*, 2002). Due to differences in units of measurements used, we

were not able to compare findings from other studies. A study directly comparing total phenolic content in tartary and common buckwheat reported higher total phenolic content (667 mg GAE/100g) in tartary buckwheat in comparison to common buckwheat seed samples (537 mg GAE/100g) (Y Liu, *et al.*, 2019). In addition, the aerial extracts were richest in total phenols followed by the stem and the seed-buckwheat extracts (A. R. Gulpinar, *et al.*, 2011; Holasova, *et al.*, 2002). Nine phenolic acids were quantified in both oat and buckwheat samples: p-hydroxybenzoic and dihydroxybenzoic acid, caffeic, p-coumaric, ferulic, vanillic, sinapic, gallic, and syringic acid (Table 1&2). The most abundant phenolic acid in oat was ferulic acid, which levels reached 149.36 mg/100g in oat grain (Kovacova & Malinova, 2007). While p-anisic (also known as 4-methoxybenzoic acid) acid was the most abundant phenolic acid in buckwheat and its levels reached 1,190 mg/100g followed by vanillic acid which content reached 311.7mg/100g (Dziedzic, *et al.*, 2018; Sytar, 2015). All polyphenolic compounds identified in oat and buckwheat are listed in Tables 1 & 2. Different solvents extraction methodology may explain differences in the extraction efficiency of phenolic compounds found in the current review. For example, Dziedzic *et al.*, found that the sum of phenolic compounds obtained from different morphological parts of buckwheat (seed, flower, root) using methanol was higher than compounds extracted using water (Dziedzic, *et al.*, 2018). Besides differences caused by solvents used, across the studies, different cultivars, genetic varies, and growing conditions were used, additionally affecting our findings. Indeed, all thermal treatments led to losses of total phenolic acid concentration in the common buckwheat samples, the greatest losses were observed with microwaving (–51.9%), followed by roasting (–33.3%), steaming (–23.5%) and boiling (–10.5%). Yet, an increase in certain phenolic acids was also noted: roasting resulted in a 3.56-fold higher concentration of syringic acid compared to the raw common buckwheat samples. Boiling and steaming caused a slight increase in p-hydroxybenzoic acid and vanillic acid+caffeic acid and p-coumaric acid+syringaldehyde, respectively (Y Liu, *et al.*, 2019).

### 3.1.3 Avenanthramides

Avenanthramides (AVAs) are a group of unique, low-molecular-weight hydroxycinnamoyl anthranilate alkaloids uniquely present in oats. AVAs, besides affecting the flavour of oat products, have been reported to improve health parameters in animal and human studies. They have antiproliferative, antioxidant, anti-inflammatory, and anti-atherogenic properties (Leonova, *et al.*, 2008; Steadman, *et al.*, 2001; R. W. Welch, 1975).

In the current review, the amount of total AVAs in oats ranged from 0.5 mg/100g to 71.85 mg/100g (Chen, *et al.*, 2018; Horbowicz & Obendorf, 1992; Hu, *et al.*, 2019; Peng, *et al.*, 2017; Tong, *et al.*, 2014).

AVAs present in more than 20 forms of which esters of 5-hydroxyanthranilic acid with p-coumaric (AVA-A), caffeic (AVA-B) and ferulic acids (AVA-C) were the most abundant (Collins, *et al.*, 1991). Total AVAs fluctuated significantly across the studies and varieties, with maximal variation of 29- fold (Chen, *et al.*, 2018). The highest AVAs content was observed in the bran and outer layer of oat kernels (its concentration decreased from the outer layer to endosperm) (Chen, *et al.*, 2018; Hitayezu, *et al.*, 2015).

The most important factors influencing the AVAs content were the genotype and harvest time (A.R. Gulpinar, *et al.*, 2011; Tong, *et al.*, 2014).

### 3.2 Tocols

Tocols (Vitamin E) are lipid-soluble compounds commonly found in vegetable oils, cereal grains (barley, oats, wheat, rye, rice) and other sources. Tocols are comprised of tocopherols (TP) and tocotrienols (T3).

There are four homologs for TP and T3:  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$ , which differ in the number and position of methyl groups on the chroman ring structure (D. M. Peterson, *et al.*, 2007). The chemical structures of the tocols determine their vitamin E activity. Different isomers of tocols exhibit vitamin E activity in the following order  $\alpha$ TP >  $\beta$ TP >  $\alpha$ T3 >  $\gamma$ TP >  $\beta$ T3 >  $\delta$ TP, with  $\alpha$ T3 having the highest antioxidant potential (Panfili, *et al.*, 2008). In the present review, tocols content was in general higher in buckwheat than in oat (5.46-55.22mg/100g in buckwheat vs 0.50-3.61 mg/100g in oat). The highest tocopherol content was reported

in whole grains and seeds. The most abundant tocopherols were alpha-tocotrienol and gamma-tocopherol in oat and buckwheat, respectively. A detailed report on tocopherols identified in oat and buckwheat can be found in Tables 1 & 2 and Supplemental Table 2.

### 3.3 Phytosterols and fatty acids

Phytosterols, plant-derived sterols, and stanols have been extensively studied due to their cholesterol-lowering effect by inhibiting the absorption of intestinal cholesterol (Leonova, *et al.*, 2008). In cereal grains, phytosterols can be found as free sterols, sterolesters, sterolglycosides and acylated sterolglycosides. Long-chain fatty acids are the major constituents of oil in grain crops and are an essential source of dietary energy.

The range of total phytosterol content was from 35mg/100g to 68.2 mg/100g in oats and 19mg/100g to 139 mg/100g in buckwheat (Table 1&2, Supplemental Table 3). The most abundant sterol in oat and buckwheat was sitosterol accounting for more than 50-70 % of total sterols, followed by campesterol. Cycloartanol was detected as a unique sterol in roasted and raw buckwheat products (Duve & White, 1991). When comparing the phytosterol composition of raw and roasted buckwheat grains, raw buckwheat grains (BG-I) had much more stigmasterol (1.3 mg/g of lipids), while the contents of campesterol, sitosterol, sitostanol, avenasterol,  $\Delta$ -7-stigmasterol, and cycloartanol were significantly lower (Hamberg, *et al.*, 1998). Raw buckwheat groats contained more campesterol (3.6 mg/g of lipids), avenasterol (3.2 mg/g of lipids) and cycloartanol (1.9 mg/g of lipids) in comparison with roasted buckwheat groat (R. W. Welch, 1975).

The fatty acid content in oat and buckwheat consisted mainly of high levels of unsaturated oleic (18:1) and linoleic (18:2), and saturated palmitic (16:0) acids and lower levels of stearic (18:0) and linoleic (18:3) acids (Supplemental Table 3). Changes in the relative abundance of fatty acids were linked to the different anatomic part of the plant, genetic variants, and influence of various environmental growth factors was reported (Doehlert, *et al.*, 2013; Sinkovic, *et al.*, 2020b; R. W. Welch, 1975),

### 3.4 Polysaccharides

Polysaccharides represent a structurally diverse class of biological macromolecules, with a wide range of physiological functions, including anticarcinogenic, hypolipidemic, immunoregulatory, antioxidant and neuroprotective activities (A.R. Gulpinar, *et al.*, 2011).  $\beta$ -glucans are heterogeneous non-starch polysaccharides, which occur in the walls of the endosperm cells (Dorrell, 1971b).  $\beta$ -glucans have been associated with increased intestinal viscosity and linked with improved health outcomes, such as improvements in serum blood lipids and enhanced efficiency of the immune system (Dorrell, 1971b). The  $\beta$ -glucan content in oat ranged from 0.03% to 8.39% in germ and endosperm. In general,  $\beta$ -glucan content was higher in oats (5,180- 28,200 mg/100g) than buckwheat (1,260-3,500 mg/100g) (Hozová, *et al.*, 2008). Factors such as temperature, pH, and pH-temperature interaction of the extraction process may influence  $\beta$ -glucan recovery and functionality (Tsuzuki, *et al.*, 1991). The amylose was the predominant polysaccharide in buckwheat, reaching 25,000 mg/100g in buckwheat starch. In addition, the levels of fagopyritols, mono-, di- and tri- galactosyl derivatives of D-chiro-inositol, were high buckwheat. Total fagopyritols content in common buckwheat ranged from 128.2 to 2774.5 mg/100g, suggesting that the common buckwheat starch may be the best source of fagopyritols.

### 3.5 Saponins

Saponins are found in various plant species, yet the oat represents the only saponin accumulating cereal. Besides being responsible for the bitter flavour of oat and playing a role in plant defense against environmental fungi attacks (Osbourn, *et al.*, 1994), saponins are suggested to promote human health by lowering cholesterol levels and affecting the immune system (Shi, *et al.*, 2004). Avenacins, stored directly in their active form, were reported in oat roots, while avenacosides (inactive forms) were reported in leaves and grains (Mary, *et al.*, 1986; Rudolf Tschesche & Lauven, 1971; Rudolf Tschesche & Wiemann,

1977). Avenacosides A (1) and B (2) are converted to their active forms: 26-desglucoavenacoside A (3) and 26-desglucoavenacoside B (4), Table 1, Supplemental Table 3. Avenacoside A represented 41.9%–60.6% and 37.1%–57.7% of total saponin content in grain and in husks respectively. Avenacoside B represented 35.8%–55.2% in grain and 13.8%–49.0% in husks of total saponin content. The content of 26-desglucoavenacoside A reached 6.4% in grain 48.3% in husks (Pecio, *et al.*, 2013). Yet, the content of avenacosides varied due to differences in cultivars, growth conditions, and sensitivity and precision of quantification methods used across different studies (Onning & Asp, 1993).

#### 4. Strengths and Limitations of Current Review

This review was conducted following the 24-Step guideline on how to perform a systematic review, and following a predesigned review protocol (Muka, *et al.*, 2020). To exhaust relevant studies in the literature, information specialists created a highly sensitive search strategy, and additional resources were searched including the reference list of included studies. Thus, in this comprehensive systematic review of the literature, we provided the first direct comparison of phytochemical content in oat and buckwheat. However, our systematic review has some limitations worth mentioning. Although a highly sensitive search strategy was used, we were not able to search all existing online databases, and we cannot exclude the possibility of missed articles. Although we did not apply any restrictions on language and publication date, we may have missed the articles published in languages other than English. Furthermore, due to the high heterogeneity among the included studies (i.e. different anatomical part of the plant used, extraction methods or reporting units), a quantification of different results was challenging and in the results section we have focused on narrative summary of the most relevant evidence.

#### 5. Implications for Future Research

The current evidence on phytochemical compositions of oat and buckwheat is highly heterogeneous (due to different genotypes, anatomical parts, cooking methods, extraction methods, etc.). A large number of studies included in our systematic review were published before 2010s (n=91, 59.09%). Thus, those extraction, purification and isolation methods of the active compounds in plants may not be as efficient as analyses done in the most recent studies. Future studies should take advantage of the separation technologies and the latest advances in plant genomics to investigate the genomic basis of the synthesis and function of phytochemicals, particularly based on recent advances in omics research.

Over the past three decades, the health benefits of oat have been well documented. Despite its potential to improve human health, buckwheat remains less often studied in clinical settings. Although the bioactive components responsible for the beneficial health effects of buckwheat remain insufficiently studied, its antioxidant potential and favorable lipid profile may be among the most important underlying factors. However, only a limited number of studies have examined the bioavailability of select phytochemicals (such as phenolic acids and polyphenols) from oats and buckwheat in humans and it remains insufficiently studied how these phytochemicals are metabolized in human bodies and influence human health. Thus, the findings from this review might help guide future research to further explore the health promoting aspects of oat and buckwheat in relation to the major phytochemicals and their metabolites, which in turn will shed light on health benefits these crops may deliver and their potential to be incorporated into human diet for optimal health.

## 6. Conclusions

Oat and buckwheat are good sources of carbohydrates, proteins, fiber, nutrients, and bioactive phytochemicals. Total phytosterols, tocopherols, flavonoids and rutin content were generally higher in

295 buckwheat,  $\beta$ -glucans were significantly higher in oat while avenanthramides and saponins were  
296 characteristically found in oat. Despite slight differences in the content of bioactive compounds, health  
297 benefits could be attributed to bioactive compounds originating from both plants, making them both an  
298 essential constituent of human diets. For individuals with coeliac disease, these plants have the potential  
299 to be a cornerstone of "gluten-free" diets due to its trace contents found in buckwheat and oats.  
300 However, with the recent evidence on the role of food processing affecting the nutritional contents of  
301 buckwheat and oat, the industry is put into a challenge in finding a proper balance of raw food  
302 transformation. Further research is needed to identify the most active forms and the bioactivity of these  
303 compounds present in oat and buckwheat and explore the health effects of the combination of nutrients  
304 and phytochemicals.  
305



306 Author's contribution

307 Study concept and design: Taulant Muka, Marija Glisic and Hua Kern.

308 Search strategy creation and online database search: Beatrice Minder.

309 Acquisition, collection, interpretation of data: Peter Francis Raguindin, Oche Adam Itodo, Jivko Stoyanov,  
310 Gordana M. Dejanovic, Magda Gamba, Eralda Asllanaj.

311 Drafting of the manuscript: Peter Francis Raguindin, Oche Adam Itodo, Marija Glisic and Hua Kern.

312 Critical revision of the manuscript for important intellectual content: Jivko Stoyanov, Gordana M.  
313 Dejanovic, Magda Gamba, Eralda Asllanaj, Beatrice Minder, Weston Bussler, Brandon Metzger, Taulant  
314 Muka.

315 Study supervision: Marija Glisic and Hua Kern.

316 All authors approved the final version of the manuscript.

317

318 Author's disclosure

319 This research was supported by Standard Process. HK, WB, BMetzger are scientists at Standard Process  
320 Nutrition Innovation Center. Other authors have nothing to disclose.

Table 1. Phytochemicals and metabolites identified in oat			
No.	Compound name	Source	Author, publication year
Phytosterols			
1	$\Delta^5$ -avenasterol	oat flour and bran; oat seed	(Maatta, <i>et al.</i> , 1999; Shewry, <i>et al.</i> , 2008)
2	$\Delta^7$ -avenasterol	oat flour and bran; oat seed	(Maatta, <i>et al.</i> , 1999; Shewry, <i>et al.</i> , 2008)
4	24-methylene cycloartanol	roots of oat seedlings	(Trojanowska, <i>et al.</i> , 2000)
3	7-hydroxymatairesinol	oat seed	(Smeds, <i>et al.</i> , 2009)
5	campesterol	oat flour and bran	(Maatta, <i>et al.</i> , 1999; Shewry, <i>et al.</i> , 2008)
6	cycloartenol	roots of oat seedlings	(Trojanowska, <i>et al.</i> , 2000)
7	flavonolignans	oat (Poaceae) herb/plant	(Wenzig, <i>et al.</i> , 2005)
8	lariciresinol	oat seed	(Smeds, <i>et al.</i> , 2009)
9	mediioresinol	oat seed	(Smeds, <i>et al.</i> , 2009)
10	metairesinol	oat seed	(Smeds, <i>et al.</i> , 2009)
11	pinioresinol	oat seed	(Smeds, <i>et al.</i> , 2009)
12	secoisolariciresinol	oat seed	(Smeds, <i>et al.</i> , 2009)
13	sitosterol	oat flour and bran; oat seed	(Maatta, <i>et al.</i> , 1999; Shewry, <i>et al.</i> , 2008) (Eichenberger Wa U, 1984)
14	stanols	oat flour and bran	(Shewry, <i>et al.</i> , 2008)
15	sterylglucosides	oat leaves	(Eichenberger Wa U, 1984)
16	stigmasterol	oat flour and bran	(Maatta, <i>et al.</i> , 1999; Shewry, <i>et al.</i> , 2008) (Eichenberger Wa U, 1984)
17	syringaresinol	oat seed	(Smeds, <i>et al.</i> , 2009)
Flavonoids			
18	4'-O-(erythro- $\beta$ -guaiacylglyceryl)	oat (Poaceae) herb/plant	(Wenzig, <i>et al.</i> , 2005)
19	4'-O-(threo- $\beta$ -guaiacylglyceryl)	oat (Poaceae) herb/plant	(Wenzig, <i>et al.</i> , 2005)
20	apigenin	oats	(Rice-Evans & Paganga, 1997)
21	kaempferol 3-O-(2'',3''-di-E-p-coumaeryl- $\alpha$ -L-rhamnoprinoside	oat bran	(WK Zhang, <i>et al.</i> , 2012)
22	kaempferol 3-O-(2''-O-E-p-coumaeryl-- $\alpha$ -L-rhamnoprinoside	oat bran	(WK Zhang, <i>et al.</i> , 2012)
23	kaempferol 3-O-(2''-O-E-p-coumaeryl-- $\beta$ -D-rhamnoprinoside	oat bran	(WK Zhang, <i>et al.</i> , 2012)
24	kaempferol 3-O-(3''-E-p-coumaeryl-- $\alpha$ -L-rhamnoprinoside	oat bran	(WK Zhang, <i>et al.</i> , 2012)
25	kaempferol 3-O- $\beta$ -D-glucopyranoside	oat bran	(WK Zhang, <i>et al.</i> , 2012)
26	linarin	oat bran	(WK Zhang, <i>et al.</i> , 2012)
27	luteolin	oats	(Collins, 1986)
28	myricitrin	oat bran	(WK Zhang, <i>et al.</i> , 2012)
29	quercetin	husked and common oat	(Kerlene, <i>et al.</i> , 2015)
30	quercitrin	oat bran	(WK Zhang, <i>et al.</i> , 2012)
31	rutin	husked oat	(Kerlene, <i>et al.</i> , 2015)
32	tilianin	oat bran	(WK Zhang, <i>et al.</i> , 2012)
33	tricin	oats	(Collins, 1986)
Polysaccharides			
34	$\beta$ -Glucan	starch; commercial oat bran; rolled oats	(Dawkins & Nnanna, 1995; Doehlert, <i>et al.</i> , 2013; Manthey, <i>et al.</i> , 1999; Regand, <i>et al.</i> , 2011; Shewry, <i>et al.</i> , 2008; Sowa & White, 1992 ; van den Broeck, <i>et al.</i> , 2015)

Phenolics			
33	2,2-diphenyl-1-picrylhydrazyl	hull and groat	(Varga, <i>et al.</i> , 2018)
34	4-hydrobenzoic acid	oat flour and bran	(Duve & White, 1991; Gallagher, <i>et al.</i> , 2010; Shewry, <i>et al.</i> , 2008; Soykan, <i>et al.</i> , 2019 ; Varga, <i>et al.</i> , 2018)
35	avenaluminic acid and its 3'-hydroxy and 3'-methoxy derivatives	oat groats and hulls	(Collins, <i>et al.</i> , 1991)
36	Caffeic acid	oat flour; hull and groat	(Calinoiu & Vodnar, 2019 ; Dimburg, <i>et al.</i> , 1993; Emmons & Peterson, 1999; Hitayezu, <i>et al.</i> , 2015; Skoglund, <i>et al.</i> , 2008; Sosulski, <i>et al.</i> , 1982; Soykan, <i>et al.</i> , 2019; Varga, <i>et al.</i> , 2018; Xing & White, 1997)
37	coumaric acid	husked oat	(Multari, <i>et al.</i> , 2018)
38	ferulic acid	oat flour; husked oat; rolled oat; hull and groat	(Calinoiu & Vodnar, 2019; Dokuyucu, <i>et al.</i> , 2003; Emmons & Peterson, 1999; Gallagher, <i>et al.</i> , 2010; Hitayezu, <i>et al.</i> , 2015; Kovacova & Malinova, 2007; Multari, <i>et al.</i> , 2018; Skoglund, <i>et al.</i> , 2008; Sosulski, <i>et al.</i> , 1982; Soykan, <i>et al.</i> , 2019; Varga, <i>et al.</i> , 2018; V. Verardo, <i>et al.</i> , 2011; Xing & White, 1997)
39	gallic acid	hull and groat	(Brindzova, <i>et al.</i> , 2008; Chen, <i>et al.</i> , 2018; Emmons & Peterson, 1999)
40	o-coumaric acid	husked oat	(Gallagher, <i>et al.</i> , 2010; Multari, <i>et al.</i> , 2018; Xing & White, 1997)
41	p-cinnamic	oat milling fractions; husked oat	(Bratt, <i>et al.</i> , 2003; Dimberg, <i>et al.</i> , 2005; Hitayezu, <i>et al.</i> , 2015; Multari, <i>et al.</i> , 2018)
42	p-coumaric acid	oat flour and bran; rolled oat; hull and groat	(Calinoiu & Vodnar, 2019; Dokuyucu, <i>et al.</i> , 2003; Emmons & Peterson, 1999; Shewry, <i>et al.</i> , 2008; Skoglund, <i>et al.</i> , 2008; Soykan, <i>et al.</i> , 2019; Varga, <i>et al.</i> , 2018; Xing & White, 1997)
43	p-hydroxybenzaldehyde	hull and groat	(Emmons & Peterson, 1999 ; Soykan, <i>et al.</i> , 2019 ; Varga, <i>et al.</i> , 2018 ; V. Verardo, <i>et al.</i> , 2011)
44	p-hydroxybenzoic acid	oat flour	(Calinoiu & Vodnar, 2019; Sosulski, <i>et al.</i> , 1982 ; Xing & White, 1997 )
45	phytic acid	oat products	(Larsson & Sandberg, 1992)
46	sinapic acid	oat flour	(Calinoiu & Vodnar, 2019 ; Shewry, <i>et al.</i> , 2008 ; Soykan, <i>et al.</i> , 2019; Xing & White, 1997 )
47	syringaldehyde	husked oat	(Gallagher, <i>et al.</i> , 2010; Multari, <i>et al.</i> , 2018 )
48	syringic	oat flour; husked oat	(Shewry, <i>et al.</i> , 2008; Sosulski, <i>et al.</i> , 1982; Soykan, <i>et al.</i> , 2019; Varga, <i>et al.</i> , 2018 ; V. Verardo, <i>et al.</i> , 2011)
49	trans-cinnamic acid	hull and groat	(Emmons & Peterson, 1999)
50	trans-ferulic	oat flour	(Sosulski, <i>et al.</i> , 1982);
51	vanillic acid	oat flour; husked oat	(Calinoiu & Vodnar, 2019; Emmons & Peterson, 1999 ; Gallagher, <i>et al.</i> , 2010 ; Garleb, <i>et al.</i> , 1991; Hitayezu, <i>et al.</i> , 2015; Multari, <i>et al.</i> , 2018 ; Shewry, <i>et al.</i> , 2008; Sosulski, <i>et al.</i> , 1982; Soykan, <i>et al.</i> , 2019; Varga, <i>et al.</i> , 2018; V. Verardo, <i>et al.</i> , 2011; Xing & White, 1997 )
Alkaloids			
52	AVA-2fd	seeds	(de Bruijn, <i>et al.</i> , 2019)

53	AVA-2pd	seeds	(de Bruijn, <i>et al.</i> , 2019)
54	AVA-5fd	seeds	(de Bruijn, <i>et al.</i> , 2019)
55	AVA-5pd	seeds	(de Bruijn, <i>et al.</i> , 2019)
56	AVA-2C	oat milling fractions; husked oat	(Bratt, <i>et al.</i> , 2003; Calinoiu & Vodnar, 2019; Chen, <i>et al.</i> , 2018; de Bruijn, <i>et al.</i> , 2019; Dimberg, <i>et al.</i> , 2005; Dokuyucu, <i>et al.</i> , 2003; Gunther-Jordanland, <i>et al.</i> , 2016; Hitayezu, <i>et al.</i> , 2015; Hu, <i>et al.</i> , 2019; Mattila, <i>et al.</i> , 2005; Multari, <i>et al.</i> , 2018; Ortiz-Robledo, <i>et al.</i> , 2013; D. M. Peterson & Dimberg, 2008; Pridal, <i>et al.</i> , 2018; Shewry, <i>et al.</i> , 2008; Skoglund, <i>et al.</i> , 2008; Varga, <i>et al.</i> , 2018; V. Verardo, <i>et al.</i> , 2011; Xie, <i>et al.</i> , 2017 )
57	AVA-2F	oat milling fractions; husked oat	(Bratt, <i>et al.</i> , 2003; Calinoiu & Vodnar, 2019; Chen, <i>et al.</i> , 2018; Dimberg, <i>et al.</i> , 2005 ; Dokuyucu, <i>et al.</i> , 2003; Gunther-Jordanland, <i>et al.</i> , 2016; Hitayezu, <i>et al.</i> , 2015; Hu, <i>et al.</i> , 2019; Mattila, <i>et al.</i> , 2005; Multari, <i>et al.</i> , 2018; Ortiz-Robledo, <i>et al.</i> , 2013; D. M. Peterson & Dimberg, 2008; Pridal, <i>et al.</i> , 2018 ; Shewry, <i>et al.</i> , 2008; Skoglund, <i>et al.</i> , 2008; Varga, <i>et al.</i> , 2018; Xie, <i>et al.</i> , 2017);
58	AVA-2P	oat milling fractions; husked oat	(Bratt, <i>et al.</i> , 2003; Calinoiu & Vodnar, 2019; Chen, <i>et al.</i> , 2018; Dimberg, <i>et al.</i> , 2005 ; Dokuyucu, <i>et al.</i> , 2003; Gunther-Jordanland, <i>et al.</i> , 2016; Hitayezu, <i>et al.</i> , 2015; Hu, <i>et al.</i> , 2019 ; Mattila, <i>et al.</i> , 2005; Multari, <i>et al.</i> , 2018; Ortiz-Robledo, <i>et al.</i> , 2013; D. M. Peterson & Dimberg, 2008; Pridal, <i>et al.</i> , 2018 ; Shewry, <i>et al.</i> , 2008; Skoglund, <i>et al.</i> , 2008; Varga, <i>et al.</i> , 2018; Xie, <i>et al.</i> , 2017 )
59	AVA A-1	oat flour and bran groat and hull	(Bratt, <i>et al.</i> , 2003; Chen, <i>et al.</i> , 2018 ; Collins, 1989 ; Gunther-Jordanland, <i>et al.</i> , 2016 ; Hu, <i>et al.</i> , 2019) (Collins, 1989; Dimburg, <i>et al.</i> , 1993)
60	AVA B	groat and hull	(Collins, 1989)
61	AVA B-1	groat and hull	(Collins, 1989)
62	AVA C	groat and hull	(Collins, 1989 ; Soycan, <i>et al.</i> , 2019)
63	AVA C-1	groat and hull	(Collins, 1989)
64	AVA D	groat and hull	(Collins, 1989)
65	AVA D-1	groat and hull	(Collins, 1989)
66	AVA E	groat and hull	(Collins, 1989)
67	AVA E-1	groat and hull	(Collins, 1989)

68	AVA-1F	oat flour	(Bratt, <i>et al.</i> , 2003; Chen, <i>et al.</i> , 2018; Collins, 1989; Gunther-Jordanland, <i>et al.</i> , 2016 )
69	AVA-1P	oat flour	(Bratt, <i>et al.</i> , 2003; Chen, <i>et al.</i> , 2018 ; Collins, 1989; Gunther-Jordanland, <i>et al.</i> , 2016 )
70	AVA-1S	oat flour	(Bratt, <i>et al.</i> , 2003; Chen, <i>et al.</i> , 2018 ; Collins, 1989; Gunther-Jordanland, <i>et al.</i> , 2016 )
71	AVA-2S	oat flour	(Gunther-Jordanland, <i>et al.</i> , 2016)
72	AVA-3F	grains	(Skoglund, <i>et al.</i> , 2008)
73	AVA-6f	seeds	(de Bruijn, <i>et al.</i> , 2019)
n.a.	avenanthramides total	oat groats, hulls and seeds; husked oat	(Bratt, <i>et al.</i> , 2003; Chen, <i>et al.</i> , 2018; Collins, <i>et al.</i> , 1991; Emmons & Peterson, 1999; Hu, <i>et al.</i> , 2019; Ishihara, <i>et al.</i> , 2014; Multari, <i>et al.</i> , 2018 ; Pridal, <i>et al.</i> , 2018; Shewry, <i>et al.</i> , 2008; Xie, <i>et al.</i> , 2017)
74	gramine	Avena sativa	(Duke, 1992)
Tocols			
75	$\alpha$ -T	various oats	(Musa Ozcan, <i>et al.</i> , 2006; D.M. Peterson & Qureshi, 1993; Shewry, <i>et al.</i> , 2008; van den Broeck, <i>et al.</i> , 2015; York, <i>et al.</i> , 1993)
76	A-T3	various oats	(Shewry, <i>et al.</i> , 2008; van den Broeck, <i>et al.</i> , 2015; York, <i>et al.</i> , 1993)
77	B-T	various oats	(Musa Ozcan, <i>et al.</i> , 2006; Shewry, <i>et al.</i> , 2008; York, <i>et al.</i> , 1993)
78	B-T3	various oats	(Shewry, <i>et al.</i> , 2008; York, <i>et al.</i> , 1993)
79	g-T	various oats	(Musa Ozcan, <i>et al.</i> , 2006; York, <i>et al.</i> , 1993)
80	6-T3	various oats	(York, <i>et al.</i> , 1993)
Folates			
81	Total folates	oat flour and bran	(Shewry, <i>et al.</i> , 2008)
Proteins/enzymes			
82	fraction I protein	Oat leaves	(Steer, <i>et al.</i> , 1968)
83	globulin	various oats	(Klose, <i>et al.</i> , 2009; Runyon, <i>et al.</i> , 2013)
84	glutelin	oat seeds/groat	(Klose, <i>et al.</i> , 2009; Runyon, <i>et al.</i> , 2013)
85	oat protein isolate	Chinese oat	(G Liu, <i>et al.</i> , 2009)
86	prolamin	oat seeds/groat	(Klose, <i>et al.</i> , 2009)
Lipids/Fatty Acids			
87	1,2-diacylglycerol	wild and cultivated oat	(Leonova, <i>et al.</i> , 2008)
88	1,3-diacylglycerol	wild and cultivated oat	(Leonova, <i>et al.</i> , 2008)
89	7-hydroxyhexadecanoic	wild and cultivated oat	(Leonova, <i>et al.</i> , 2008)
90	cylphosphatidylglycerols	oat extract	(Holmback, <i>et al.</i> , 2001)
91	1-[(9'Z),(12'Z)-octadecadienoyl]-2-[(15'R)-(C Chen, <i>et al.</i> , 2020)-(9"Z),(12"Z)-octadecadienoyl]-3-( $\alpha$ -D-galactopyranosyl-1-6- $\beta$ -D-galactopyranosyl)-glycerol.	oat seeds	(Hamberg, <i>et al.</i> , 1998)
92	galactolipid estolides	oat kernels	(Moreau, <i>et al.</i> , 2008)
93	N-acylphosphatidylethanolamines	oat extract	(Holmback, <i>et al.</i> , 2001)
94	sterylglucosides	oat leaves	(Eichenberger Wa U, 1984)

95	TAG1	wild and cultivated oat	(Leonova, <i>et al.</i> , 2008)
96	TAG2	wild and cultivated oat	(Leonova, <i>et al.</i> , 2008)
97	C:16	Oat various cultivars	(Banas, <i>et al.</i> , 2007; Brindzova, <i>et al.</i> , 2008; Dimberg, <i>et al.</i> , 2005; Duve & White, 1991; van den Broeck, <i>et al.</i> , 2015; R.W. Welch, 1975)
98	C18:0	oat various cultivars	(Banas, <i>et al.</i> , 2007; Brindzova, <i>et al.</i> , 2008; Dimberg, <i>et al.</i> , 2005; Duve & White, 1991; van den Broeck, <i>et al.</i> , 2015; R.W. Welch, 1975)
99	C18:1	oat various cultivars	(Banas, <i>et al.</i> , 2007; Brindzova, <i>et al.</i> , 2008; Dimberg, <i>et al.</i> , 2005; Duve & White, 1991; van den Broeck, <i>et al.</i> , 2015; R.W. Welch, 1975)
100	C18:2	oat various cultivars	(Banas, <i>et al.</i> , 2007; Brindzova, <i>et al.</i> , 2008; Dimberg, <i>et al.</i> , 2005; Duve & White, 1991; van den Broeck, <i>et al.</i> , 2015; R.W. Welch, 1975)
101	Avenoleic acid	oat seeds;	(Hamberg, <i>et al.</i> , 1998)
102	15-hydroxy 18:2 <sup>Δ9,12</sup> (avenoleic acid)	wild and cultivated oat	(Leonova, <i>et al.</i> , 2008)
<b>Saponins</b>			
103	3-(O-α-L-rhamnopyranosyl(1→2)-[β-D-glucopyranosyl(1→4)]-β-D-glucopyranosid)-26-O-β-D-glucopyranosyl-(25R)-furost-5-ene-3β,22,26-triol	oat flour	(Gunther-Jordanland, <i>et al.</i> , 2016)
104	3-(O-α-L-rhamnopyranosyl(1→2)-[β-D-glucopyranosyl(1→3)-β-D-glucopyranosyl(1→4)]-β-D-glucopyranosid)-26-O-β-D-glucopyranosyl-(25R)-furost-5-ene-3β,22,26-triol	oat flour	(Gunther-Jordanland, <i>et al.</i> , 2016)
105	avenacoside A (nuatigenin-3-O-(α-L-rhamnopyranosyl-(1→2)-[β-D-glucopyranosyl-(1→4)]-β-D-glucopyranoside)-26-O-β-D-glucopyranoside)	oat flour, grain	(Bahraminejad, <i>et al.</i> , 2008; Gunther-Jordanland, <i>et al.</i> , 2016; Onning & Asp, 1993; Pecio, <i>et al.</i> , 2013; Rudolf Tschesche, <i>et al.</i> , 1969; Rudolf Tschesche & Wiemann, 1977)
106	avenacoside B (nuatigenin-3-O-(α-L-rhamnopyranosyl-(1→2)-[β-D-glucopyranosyl-(1→3)-β-D-glucopyranosyl-(1→4)]-β-D-glucopyranoside)-26-O-β-D-glucopyranoside)	oat flour, grains	(Bahraminejad, <i>et al.</i> , 2008; Gunther-Jordanland, <i>et al.</i> , 2016; Onning & Asp, 1993; Pecio, <i>et al.</i> , 2013; Rudolf Tschesche & Lauven, 1971; Rudolf Tschesche & Wiemann, 1977; R. Tschesche & Wulff, 1973)
107	Avenacin A-1*		(Mary, <i>et al.</i> , 1986)
108	Avenacin A-2		(Mary, <i>et al.</i> , 1986)
109	Avenacin B-1		(Mary, <i>et al.</i> , 1986)
110	Avenacin B-2		(Mary, <i>et al.</i> , 1986)
111	26-desglucoavenacoside A	grain	(Bahraminejad, <i>et al.</i> , 2008; Pecio, <i>et al.</i> , 2013)
112	26-desglucoavenacoside B	Shoots	(Bahraminejad, <i>et al.</i> , 2008)
113	avenacins (b-Amyrin, squalene)	roots of oat seedlings	(Trojanowska, <i>et al.</i> , 2000)
* A-1, A-2, B-1 and B-2: these are trisaccharide-bearing triterpenes esterified (A-1, B-1) with N-methylantranilic acid or (A-2, B-2) benzoic acid.			




Table 2 Phytochemicals reported in buckwheat			
No.	Compound name	Source	Author, publication year
<b>Tannins</b>			
1	procyanidin B-1	Fd	(KJ Wang, <i>et al.</i> , 2005)
2	procyanidin B-2	Fd	(KJ Wang, <i>et al.</i> , 2005)
3	3,3-di-O-galloyl-procyanidinB-2	Fd	(KJ Wang, <i>et al.</i> , 2005)
4	3-O-galloyl-procyanidinB-2	Fd	(KJ Wang, <i>et al.</i> , 2005)
<b>Cyclitol</b>			
5	fagopyritol A1	Fe	(Ralph L. Obendorf, <i>et al.</i> , 2000)
6	fagopyritol A2	Fe	(Steadmana, <i>et al.</i> , 2001)
7	fagopyritol A3	Fe	(Steadmana, <i>et al.</i> , 2001)
8	fagopyritol B1	Fe	(Ralph L. Obendorf, <i>et al.</i> , 2000)
9	fagopyritol B2	Fe	(R. L. Obendorf, 1997)
10	fagopyritol B3	Fe	(R. L. Obendorf, 1997)
<b>Triterpenoids</b>			
11	ursolic acid	Ft	(JM Lee, <i>et al.</i> , 2013)
12	olean-12-en-3-ol	Fe	(F Zheng, <i>et al.</i> , 2004)
13	urs-12-en-3-ol	Fe	(F Zheng, <i>et al.</i> , 2004)
14	glutinone	Fd	(Shao, <i>et al.</i> , 2005)
15	glutanol	Fd	(Shao, <i>et al.</i> , 2005)
<b>Steroids</b>			
16	$\beta$ -sitosterol	Fd	(Bao, <i>et al.</i> , 2003)
17	daucoesterol	Fd	(Bao, <i>et al.</i> , 2003)
18	peroxidize-ergosterol	Fd	(Bao, <i>et al.</i> , 2003)
19	stigma-4-en -3,6-dione	Fd	(Bao, <i>et al.</i> , 2003)
20	$\beta$ -sitosterol-palmitate	Fd	(Bao, <i>et al.</i> , 2003)
21	6-hydroxystigmasta-4,22-dien-3-one	Fe	(F Zheng, <i>et al.</i> , 2004)
22	23S-methylcholesterol	Fe	(F Zheng, <i>et al.</i> , 2004)
23	trans-stigmast-5,22-dien-3-ol	Fe	(F Zheng, <i>et al.</i> , 2004)
24	stigmast-5,24-dien-3-ol	Fe	(F Zheng, <i>et al.</i> , 2004)
25	stigmast-5-en-3-ol	Fe	(F Zheng, <i>et al.</i> , 2004)
<b>Fatty acids</b>			
26	C24:0	Fe	(Krumina-Zemture & Beitane, 2018; Tien, <i>et al.</i> , 2018; Tsuzuki, <i>et al.</i> , 1991)
27	C24:1	Fe	(Krumina-Zemture & Beitane, 2018)
28	C12:0	Ft	(Tien, <i>et al.</i> , 2018)
29	C14:0	Ft, Fe	(Golijan, <i>et al.</i> , 2019; Krumina-Zemture & Beitane, 2018; Tien, <i>et al.</i> , 2018)
30	C15:0	Ft	(Krumina-Zemture & Beitane, 2018; Tien, <i>et al.</i> , 2018)
31	C16:0	Fe, Ft	(Dziadek, <i>et al.</i> , 2016; Golijan, <i>et al.</i> , 2019; A.R. Gulpinar, <i>et al.</i> , 2011; Khalass, <i>et al.</i> , 2018; Krumina-Zemture & Beitane, 2018; Sinkovic, <i>et al.</i> , 2020a; Tien, <i>et al.</i> , 2018 ; Tsuzuki, <i>et al.</i> , 1991)
32	C16:1	Ft	(Golijan, <i>et al.</i> , 2019; Tien, <i>et al.</i> , 2018)
33	C17:1	Ft	(Tien, <i>et al.</i> , 2018)
34	C18:0	Fe, Ft	(Dziadek, <i>et al.</i> , 2016; Golijan, <i>et al.</i> , 2019 ; A.R. Gulpinar, <i>et al.</i> , 2011; Khalass, <i>et al.</i> , 2018; Krumina-Zemture & Beitane, 2018 ; Sinkovic, <i>et al.</i> , 2020a; Tien, <i>et al.</i> , 2018; Tsuzuki, <i>et al.</i> , 1991)
35	C18:1	Fe, Ft	(Dorrell, 1971a; Dziadek, <i>et al.</i> , 2016; Golijan, <i>et al.</i> , 2019; A.R. Gulpinar, <i>et al.</i> , 2011 ; Khalass, <i>et al.</i> , 2018; Krumina-Zemture & Beitane, 2018; Sinkovic, <i>et al.</i> , 2020a; Tien, <i>et al.</i> , 2018; Tsuzuki, <i>et al.</i> , 1991)
36	C18:2	Fe, Ft	(Dziadek, <i>et al.</i> , 2016; Golijan, <i>et al.</i> , 2019; A.R. Gulpinar, <i>et al.</i> , 2011; Khalass, <i>et al.</i> , 2018; Sinkovic, <i>et al.</i> , 2020a; Tien, <i>et al.</i> , 2018; Tsuzuki, <i>et al.</i> , 1991)
37	C18:3	Fe, Ft	(Golijan, <i>et al.</i> , 2019; A.R. Gulpinar, <i>et al.</i> , 2011; Krumina-Zemture & Beitane, 2018 ; Sinkovic, <i>et al.</i> , 2020a; Tien, <i>et al.</i> , 2018; Tsuzuki, <i>et al.</i> , 1991);



38	C20:0	Fe, Ft	(Golijan, <i>et al.</i> , 2019; A.R. Gulpinar, <i>et al.</i> , 2011; Sinkovic, <i>et al.</i> , 2020a ; Tsuzuki, <i>et al.</i> , 1991);
39	C20:1	Fe, Ft	(Golijan, <i>et al.</i> , 2019; A.R. Gulpinar, <i>et al.</i> , 2011; Krumina-Zemture & Beitane, 2018; Tien, <i>et al.</i> , 2018; Tsuzuki, <i>et al.</i> , 1991)
40	C20:5	Fe	(Krumina-Zemture & Beitane, 2018)
41	C22:0	Fe, Ft	(Tien, <i>et al.</i> , 2018)
42	C22:1	Fe	(A.R. Gulpinar, <i>et al.</i> , 2011; Tien, <i>et al.</i> , 2018; Tsuzuki, <i>et al.</i> , 1991)
43	C23:0	Fe	(Krumina-Zemture & Beitane, 2018)
44	4,7-dihydroxy-3,7-dimethyl-octa-2(E),5(E)-dienoic acid	Fe	(Cho, <i>et al.</i> , 2006)
45	6,7-dihydroxy-3,7-dimethyl-octa-2(Z),4(E)-dienoic acid	Fe	(Cho, <i>et al.</i> , 2006)
46	6,7-dihydroxy-3,7-dimethyl-octa-2(E),4(E)-dienoic acid	Fe	(Cho, <i>et al.</i> , 2006)
<b>Polysaccharides</b>			
40	amylopectin	Fe	(Qin, <i>et al.</i> , 2010; Takahama & Hirota, 2010)
47	amylose	Fe	(Qin, <i>et al.</i> , 2010; Takahama & Hirota, 2010)
41	BBF1-3	Fl	(Lin & Lin, 2016)
48	BWPSs	Fe	(Zemnukhova, <i>et al.</i> , 2007; Zemnukhova, <i>et al.</i> , 2004 )
49	CBF1-3	Fe	(Lin & Lin, 2016)
39	galactinol	Fe	(Steadman, <i>et al.</i> , 2000)
50	inositol	Fe	(Steadman, <i>et al.</i> , 2001)
51	myo-inositol	Fe	(Steadman, <i>et al.</i> , 2000)
52	SDF (carboxymethylated)	Fe	(KY Lee, <i>et al.</i> , 2014)
53	SDF (methanolysis)	Fe	(Wefers & Bunzel, 2015)
54	sucrose	Fe	(Steadman, <i>et al.</i> , 2000)
55	TBP-1-3	Fe	(Yan, <i>et al.</i> , 2011)
56	TBP-II	Ft	(XT Wang, <i>et al.</i> , 2016)
57	$\alpha$ -galactosides	Fe	(Steadman, <i>et al.</i> , 2000)
	Beta-glucan	Fe	(Hozová, <i>et al.</i> , 2008)
<b>Phenols</b>			
58	1,3,6-tri-feruloyl-6-p-coumaroyl sucrose	Ft	(Ren, <i>et al.</i> , 2013)
59	1,3,6-tri-p-coumaroyl-6-feruloyl sucrose	Ft	(Ren, <i>et al.</i> , 2013)
60	1,3-dimethoxy-2-O-b-xylopyranosyl-5-O-glucopyranosyl-benzene	Fd	(Bai, <i>et al.</i> , 2007)
61	3,4-dihydroxybenzaldehyde	Fd, Fe	(Shao, <i>et al.</i> , 2005); (Watanabe, <i>et al.</i> , 1997)
62	3,6-di-p-coumaroyl-1,6-di-feruloyl sucrose	Ft	(Ren, <i>et al.</i> , 2013)
63	benzoic acid	<b>Fd</b>	(Park, <i>et al.</i> , 2019; Wu, <i>et al.</i> , 2008)
64	caffeic acid	<b>Ft, Fe</b>	(Alvarez-Jubete, <i>et al.</i> , 2010; Park, <i>et al.</i> , 2019; Terpin, <i>et al.</i> , 2016; Wefers & Bunzel, 2015; Xu, <i>et al.</i> , 2002)
65	chlorogenic acid	Fe, Ft	(SJ Kim, <i>et al.</i> , 2008; Kraujaliene, <i>et al.</i> , 2017; Y Liu, <i>et al.</i> , 2019; Pankaja, 2011; Park, <i>et</i>

			<i>al.</i> , 2019; Sadauskiene, <i>et al.</i> , 2018; Sytar, 2015)
66	Neochloreogenic	Fe	(Sadauskiene, <i>et al.</i> , 2018)
67	Fagopyrin	Fe	(Habtemariam, 2019; Sadauskiene, <i>et al.</i> , 2018 )
68	Tannic acid	Fe	(Sadauskiene, <i>et al.</i> , 2018)
69	Hyperoside	Fe	(Quettier-Deleu, <i>et al.</i> , 2000; Sadauskiene, <i>et al.</i> , 2018)
70	ferulic acid (tras- and cis- isomers)	Ft	(Hung & Morita, 2008; Y Liu, <i>et al.</i> , 2019; Sedej, <i>et al.</i> , 2012; Wefers & Bunzel, 2015; Xu, <i>et al.</i> , 2002)
71	gallic acid	Fd, Fe	(F Li, <i>et al.</i> , 2013; Y Liu, <i>et al.</i> , 2019; Park, <i>et al.</i> , 2019; Sytar, 2015; Terpinc, <i>et al.</i> , 2016; KJ Wang, <i>et al.</i> , 2005)
72	p-coumaric acid	Ft	(Sytar, 2015; Wefers & Bunzel, 2015; Xu, <i>et al.</i> , 2002)
73	p-hydroxybenzoic acid	Fd	(Y Liu, <i>et al.</i> , 2019; Tian & Xu, 1997)
74	Protocatechuic acid	Ft	(Y Liu, <i>et al.</i> , 2019; Sedej, <i>et al.</i> , 2012; Watanabe, <i>et al.</i> , 1997)
75	protocatechuic acid methyl ester	Fd	(Shao, <i>et al.</i> , 2005)
76	resveratrol	Fe	(Watanabe, <i>et al.</i> , 1997)
77	Sinapic acid	Ft	(Y Liu, <i>et al.</i> , 2019; Sedej, <i>et al.</i> , 2012; Wefers & Bunzel, 2015)
78	syringic acid	Ft	(XD Guo, <i>et al.</i> , 2011; Y Liu, <i>et al.</i> , 2019; Xu, <i>et al.</i> , 2002)
79	taroside (1,3,6,61-tetra-feruloyl sucrose)	Ft	(Ren, <i>et al.</i> , 2013)
80	tatariside A-G	Ft	(C Zheng, <i>et al.</i> , 2012)
81	vanillic acid	Ft	(Sedej, <i>et al.</i> , 2012; Xu, <i>et al.</i> , 2002)
82	Caffeic acid hexose	Fe	(Vito Verardo, <i>et al.</i> , 2010)
83	2-Hydroxy-3-O-b-Dglucopyranosil-benzoic acid	Fe	(Vito Verardo, <i>et al.</i> , 2010)
84	Methoxy-cinnamic acid	Fe, Ft	(Sytar, 2015)
85	Salicylic acid	Fe, Ft	(Sytar, 2015)
86	p-Anisic acid	Fe, Ft	(Sytar, 2015)
<b>Flavonoids</b>			
87	(-)-epicatechin	Fe, Fd	(Kalinova, <i>et al.</i> , 2006; Park, <i>et al.</i> , 2019; Quettier-Deleu, <i>et al.</i> , 2000; Watanabe, <i>et al.</i> , 1997)
88	(-)-epicatechin-3-O-(3,4-di-O-methyl)-gallate	Fe	(Vito Verardo, <i>et al.</i> , 2010; Watanabe, <i>et al.</i> , 1997)
89	(-)-epicatechin-3-O-p-hydroxybenzoate	Fe	(Watanabe, <i>et al.</i> , 1997)
90	(+)-catechin	Fd	(Park, <i>et al.</i> , 2019; KJ Wang, <i>et al.</i> , 2005)
91	(+)-catechin-7-O-glucoside		(Watanabe, <i>et al.</i> , 1997)
92	(3-methoxyphenyl)-2-piperidinemethanol	Fd	(Shao, <i>et al.</i> , 2005)
93	(Epi)afzelchine(epi)catechin isomer A	Fe	(Vito Verardo, <i>et al.</i> , 2010)
94	(Epi)afzelchine(Epi)catechin isomer B	Fe	(Vito Verardo, <i>et al.</i> , 2010)
95	(Epi)afzelchine(epi)catechin isomer C	Fe	(Vito Verardo, <i>et al.</i> , 2010)
96	(Epi)afzelchine(epi)catechin isomer D	Fe	(Vito Verardo, <i>et al.</i> , 2010)
97	1-O-Caffeoyl-6-O-alpha-rhamnopyranosyl-beta-glucopyranoside (swertiamacroside)	Fe	(Vito Verardo, <i>et al.</i> , 2010)
98	3, 4-dihydroxy benzamine	Fd	(Shao, <i>et al.</i> , 2005)
99	3,5-dimethylquercetin	Fd	(KJ Wang, <i>et al.</i> , 2005)
100	3',4'-methylenedioxy-7-hydroxy-6-isopentenyl flavone	Fd	(Saxena, 1987)
101	3-methylgossypetin-8-O-D-glucopyranoside	Fd	(KJ Wang, <i>et al.</i> , 2005)
102	3-methylquercetin	Fd	(KJ Wang, <i>et al.</i> , 2005)
103	5, 5'-di-α-furaldehyde dimethyl ester	Ft	(Tian & Xu, 1997)
104	5,7,31,41-tetramethylquercetine-3-O-rutinoside	Ft	(Saxena, 1987)
105	7-hydroxycoumarin	Ft	Sun, B.H et al, 2008

106	Apigenin	Fe	(Terpinc, <i>et al.</i> , 2016)
107	aromadendrin-3-O-D-galactoside		(Watanabe, <i>et al.</i> , 1997)
108	Catechin-glucoside	Fe	(Vito Verardo, <i>et al.</i> , 2010)
109	emodin	Fd	(Bao, <i>et al.</i> , 2003)
110	emodin-8-O- $\beta$ -D-glucopyranoside	Fd	Wang, K. <i>et al.</i> , 2005
111	Epiafzelchine epicatechin-O-methyl gallate	Fe	(Vito Verardo, <i>et al.</i> , 2010)
112	Epiafzelchine-epicatechin-O-di methyl gallate	Fe	(Vito Verardo, <i>et al.</i> , 2010)
113	Epicatechin-gallate	Fe	(Quettier-Deleu, <i>et al.</i> , 2000; Vito Verardo, <i>et al.</i> , 2010)
114	fagopyrin	Fe, Ft	(Habtemariam, 2019; Kreft, <i>et al.</i> , 2013)
115	fructose	Ft	(KJ Wang, <i>et al.</i> , 2005)
116	hesperidin	Fd	(Wu, <i>et al.</i> , 2008)
117	hyperin/isoquercitrin (quercetin-3-O-glucoside)	Fe	(Ren, <i>et al.</i> , 2013; Watanabe, <i>et al.</i> , 1997)
118	isoorientin	Fe, Ft	(T Chen, <i>et al.</i> , 2020; HJ Kim, <i>et al.</i> , 2011; SJ Kim, <i>et al.</i> , 2008; Kraujaliene, <i>et al.</i> , 2017; Nam, <i>et al.</i> , 2015; Pankaja, 2011)
119	isovitexin	Fe, Ft	(T Chen, <i>et al.</i> , 2020; Dietrych-Szostak & Oleszek, 1999; SJ Kim, <i>et al.</i> , 2008; Nam, <i>et al.</i> , 2015; Pankaja, 2011)
120	kaempferol	Fd, Ft	(Jiang, <i>et al.</i> , 2015; JM Lee, <i>et al.</i> , 2013; Terpinc, <i>et al.</i> , 2016)
121	kaempferol-3-O-galactoside	Ft	(Bao, <i>et al.</i> , 2003; Jiang, <i>et al.</i> , 2015; KJ Wang, <i>et al.</i> , 2005)
122	kaempferol-3-O-glucoside	Ft	(Bao, <i>et al.</i> , 2003; Jiang, <i>et al.</i> , 2015; KJ Wang, <i>et al.</i> , 2005)
123	kaempferol-3-O-glucoside-7-O-glucoside	Fe	(Watanabe, <i>et al.</i> , 1997)
124	kaempferol-3-O-rutinoside	Ft	(Bao, <i>et al.</i> , 2003; Jiang, <i>et al.</i> , 2015; KJ Wang, <i>et al.</i> , 2005)
125	kaempferol-3-O-sophoroside	Fe	(Watanabe, <i>et al.</i> , 1997)
126	luteolin	Fd	(Shao, <i>et al.</i> , 2005; Terpinc, <i>et al.</i> , 2016)
127	Luteolin-glycosid	Fe	(Vito Verardo, <i>et al.</i> , 2010)
128	myricetin	Fe, Fd	(Kalinova, <i>et al.</i> , 2006)
129	naringenin	Fe	(Pankaja, 2011; Terpinc, <i>et al.</i> , 2016)
130	n-butyl- $\beta$ -D-fructopyranoside	Fd	(Shao, <i>et al.</i> , 2005)
131	N-trans-feruloyltyramine	Ft	(Ren, <i>et al.</i> , 2013)
132	orientin	Fe, Ft	(T Chen, <i>et al.</i> , 2020; HJ Kim, <i>et al.</i> , 2011; SJ Kim, <i>et al.</i> , 2008; Nam, <i>et al.</i> , 2015)
133	Procyanidin B2	Fe	(Vito Verardo, <i>et al.</i> , 2010); (Quettier-Deleu, <i>et al.</i> , 2000)
134	Procyanidin B2 dimethyl gallate	Fe	(Vito Verardo, <i>et al.</i> , 2010)
135	Procyanidin B2-3-O-gallate	Fe	(Vito Verardo, <i>et al.</i> , 2010)
136	Quercetin	Ft	(T Chen, <i>et al.</i> , 2020 ; Ge & Wang, 2020; Kočevr Glavač, <i>et al.</i> , 2017; JM Lee, <i>et al.</i> , 2013; Sadauskiene, <i>et al.</i> , 2018; Sedej, <i>et al.</i> , 2012; Terpinc, <i>et al.</i> , 2016; Vito Verardo, <i>et al.</i> , 2010)
137	quercetin-3-O-(211-O-p-hydroxy-coumaroyl)-glucoside	Fd	(KJ Wang, <i>et al.</i> , 2005);
138	quercetin-3-O-[ $\beta$ -D-xyloxy-(1 $\rightarrow$ 2)- $\beta$ -L-rhamnoside]	Ft	(Ren, <i>et al.</i> , 2013)
139	quercetin-3-O- $\beta$ -D-galactoside	Ft, Fe	(Nam, <i>et al.</i> , 2015; Ren, <i>et al.</i> , 2013; Watanabe, <i>et al.</i> , 1997)
140	quercetin-3-O-rutinoside-31-O- $\beta$ -glucopyranoside	Fd, Ft	(Wu, <i>et al.</i> , 2008)
141	quercetin-3-O-rutinoside-7-O-galactoside	Ft	(Saxena, 1987)
142	quercitrin (quercetin-3-O-rhamnoside)	Fd, Ft	(Ren, <i>et al.</i> , 2013; KJ Wang, <i>et al.</i> , 2005)
143	rhamnetin	Fd	(Wu, <i>et al.</i> , 2008)
144	rutin	Fd, Fe, Ft	(T Chen, <i>et al.</i> , 2020 ; Dietrych-Szostak & Oleszek, 1999; Fabjan, <i>et al.</i> , 2003; Ge & Wang, 2020 ; XD Guo, <i>et al.</i> , 2011; Habtemariam, 2019; Holasovaa, <i>et al.</i> , 2002; Hung & Morita, 2008; Kalinova, <i>et al.</i> , 2006; Keriene, <i>et al.</i> , 2015 ; HJ Kim, <i>et al.</i> , 2011; Kočevr Glavač, <i>et al.</i> , 2017; Kraujaliene, <i>et al.</i> , 2017; Kreft, <i>et al.</i> , 1999; Nam, <i>et al.</i> , 2015; B. Dave Oomah & Giuseppe Mazza, 1996; Park, <i>et al.</i> , 2019; Quettier-Deleu, <i>et al.</i> , 2000; Sadauskiene, <i>et al.</i> , 2018; Sedej, <i>et al.</i> , 2012; KJ Wang, <i>et al.</i> , 2005)

145	squalene	Fe	(F Zheng, <i>et al.</i> , 2004)
146	succinic acid	Fd	(Tian & Xu, 1997)
147	sucrose	Ft	(KJ Wang, <i>et al.</i> , 2005)
148	taxifolin-3-O-D-xyloside	Fe	(Watanabe, <i>et al.</i> , 1997)
149	uracil	Ft	(Bao, <i>et al.</i> , 2003)
150	vitexin	Fe, Ft	(T Chen, <i>et al.</i> , 2020; HJ Kim, <i>et al.</i> , 2011; SJ Kim, <i>et al.</i> , 2008; Nam, <i>et al.</i> , 2015; Pankaja, 2011)
<b>Vitamins</b>			
151	A ( $\beta$ -carotene)	Fe, Ft,Fd	(Gabrovská, <i>et al.</i> , 2002)
152	B1 (thiamine)	Fe, Ft,Fd	(Bonafaccia, <i>et al.</i> , 2003)
153	B2 (riboflavin)	Fe, Ft,Fd	(Bonafaccia, <i>et al.</i> , 2003)
154	B5 (pantothenic acid)	Fe, Ft,Fd	(Gabrovská, <i>et al.</i> , 2002)
155	B6 (pyridoxine)	Fe, Ft,Fd	(Bonafaccia, <i>et al.</i> , 2003)
156	C (ascorbic acid)	Fe, Ft,Fd	(Lintschinger, <i>et al.</i> , 1997)
157	E (tocopherols)	Fe, Ft,Fd	(SL Kim, Kim, S. K., and Park, C. H., 2002)
158	Alpha-tocopherol	Fe	(Sedej, <i>et al.</i> , 2012)
159	Beta--tocopherol	Fe	(Sedej, <i>et al.</i> , 2012)
160	Gamma-tocopherol	Fe	(Sedej, <i>et al.</i> , 2012; F Zheng, <i>et al.</i> , 2004)
<b>Minerals</b>			
161	Boron	Fe	(Ikeda, <i>et al.</i> , 2006; Steadman, <i>et al.</i> , 2001)
162	Calcium	Fe, Ft,Fd	(Ikeda, <i>et al.</i> , 2005; Q Zhang & Xu, 2017)
163	Copper	Fe, Ft,Fd	(Ikeda, <i>et al.</i> , 2005; Q Zhang & Xu, 2017)
164	Iron	Fe, Ft,Fd	(Ikeda, <i>et al.</i> , 2002; Q Zhang & Xu, 2017)
165	Magnesium	Fe, Ft,Fd	(Ikeda, <i>et al.</i> , 2005)
166	Manganese	Fe, Ft,Fd	(Ikeda, <i>et al.</i> , 2005; Q Zhang & Xu, 2017)
167	Phosphorus	Fe, Ft,Fd	(Ikeda, <i>et al.</i> , 2005; Q Zhang & Xu, 2017)
168	Potassium	Fe, Ft,Fd	(Ikeda, <i>et al.</i> , 2005)
169	Zinc	Fe, Ft,Fd	(Ikeda, <i>et al.</i> , 2005; Q Zhang & Xu, 2017)
<b>Protein</b>			
170	Albumin	Ft	(X Guo, <i>et al.</i> , 2007; Siwatch, <i>et al.</i> , 2019)
171	Globulin	Ft	(X Guo, <i>et al.</i> , 2007)
172	Prolamin	Ft	(X Guo, <i>et al.</i> , 2007)
173	Glutelin	Ft	(X Guo, <i>et al.</i> , 2007)
174	TBPC or TBP	Ft	(C Zhang, <i>et al.</i> , 2005; Z Zhang, <i>et al.</i> , 1999)
175	TBWSP31	Ft	(X Guo, <i>et al.</i> , 2007)
176	TBTI	Ft	(Z Wang, <i>et al.</i> , 2004)
177	DTPF	Ft	(CH Li, <i>et al.</i> , 2002)
178	TBa, TBb, TBt	Ft	(Z Wang, <i>et al.</i> , 2006)

322 **References**

- 323 Ahmed, A., Khalid, N., Ahmad, A., Abbasi, N. A., Latif, M. S. Z., & Randhawa, M. A. (2014).  
 324 Phytochemicals and biofunctional properties of buckwheat: a review. *Journal of Agricultural*  
 325 *Science*, 152(3), 349-369.
- 326 Alvarez-Jubete, L., Wijngaard, H., Arendt, E. K., & Gallagher, E. (2010). Polyphenol composition and in  
 327 vitro antioxidant activity of amaranth, quinoa buckwheat and wheat as affected by sprouting  
 328 and baking. *Food Chemistry*, 119(2), 770-778.
- 329 Arendt, E. K., & Zannini, E. (2013). Buckwheat. In *Cereal Grains for the Food and Beverage Industries*,  
 330 (pp. 369-408).
- 331 Bahraminejad, S., Asenstorfer, R. E., Riley, I. T., & Schultz, C. J. (2008). Analysis of the Antimicrobial  
 332 Activity of Flavonoids and Saponins Isolated from the Shoots of Oats (*Avena sativa* L.).  
 333 *Journal of Phytopathology*, 156, 1-7.
- 334 Bai, Z. Z., Zhang, X. H., Xuan, L. J., & Mo, F. K. (2007). A phenolic glycoside from *Fagopyrum dibotrys*  
 335 (D. Don) Hara. *Chinese Chemical Letters*, 18(9), 1087-1088.
- 336 Banas, A., Debski, H., Banas, W., Heneen, W. K., Dahlqvist, A., Bafor, M., Gummeson, P. O., Marttila,  
 337 S., Ekman, A., Carlsson, A. S., & Stymne, S. (2007). Lipids in grain tissues of oat (*Avena sativa*):  
 338 differences in content, time of deposition, and fatty acid composition. *Journal of*  
 339 *Experimental Botany*, 58(10), 2463-2470.
- 340 Bao, T. N., Peng, S. L., Zhou, Z. Z., & Li, B. G. (2003). Chemical constituents of *Fagopyrum tataricum*  
 341 (Linn) Gaertn. *Natural Products Research and Development*, 15, 24-26.
- 342 Bonafaccia, G., Gambelli, L., Fabjan, N., & Kreft, I. (2003). Trace elements in flour and bran from  
 343 common and tartary buckwheat. *Food Chemistry*, 83(1), 1-5.
- 344 Borneo, R., & Leon, A. E. (2012). Whole grain cereals: functional components and health benefits.  
 345 *Food & Function*, 3(2), 110-119.
- 346 Boz, H. (2015). Phenolic Amides (Avenanthramides) in Oats – A review. *Czech Journal of Food*  
 347 *Sciences*, 33(5), 399-404.
- 348 Bratt, K., Sunnerheim, K., Bryngelsson, S., Fagerlund, A., Engman, L., Andersson, R. E., & Dimberg, L.  
 349 H. (2003). Avenanthramides in oats (*Avena sativa* L.) and structure-antioxidant activity  
 350 relationships. *Journal of Agricultural and Food Chemistry*, 51(3), 594-600.
- 351 Brindzova, L., Certik, M., Raptá, P., Zalibera, M., Mikulajova, A., & Takacsova, M. (2008). Antioxidant  
 352 activity, beta-glucan and lipid contents of oat varieties. *Czech Journal of Food Sciences*, 26(3),  
 353 163-173.
- 354 Calinoiu, L. F., & Vodnar, D. C. (2019). Thermal Processing for the Release of Phenolic Compounds  
 355 from Wheat and Oat Bran. *Biomolecules*, 10(1), 21.
- 356 Chen, C., Li, T., Chen, Z., Wang, L., & Luo, X. (2020). Absorption Rates and Mechanisms of  
 357 Avenanthramides in a Caco-2 Cell Model and Their Antioxidant Activity during Absorption.  
 358 *Journal of Agricultural and Food Chemistry*, 68(8), 2347-2356.
- 359 Chen, C., Wang, L., Wang, R., Luo, X., Li, Y., Li, J., Li, Y., & Chen, Z. (2018). Phenolic contents, cellular  
 360 antioxidant activity and antiproliferative capacity of different varieties of oats. *Food*  
 361 *Chemistry*, 239, 260-267.
- 362 Chen, T., Piao, M., Ehsanur Rahman, S. M., Zhang, L., & Deng, Y. (2020). Influence of fermentation on  
 363 antioxidant and hypolipidemic properties of maifanite mineral water-cultured common  
 364 buckwheat sprouts. *Food Chemistry*, 321, 126741.
- 365 Cho, J. Y., Moon, J. H., Kim, H. K., Ma, S. J., Kim, S. J., Jang, M. Y., Kawazoe, K., Takaishi, Y., & Park, K.  
 366 H. (2006). Isolation and structural elucidation of antimicrobial compounds from buckwheat  
 367 hull. *Journal of Microbiology and Biotechnology*, 16(4), 538-542.

- Collins, F. W. (1989). Oat Phenolics - Avenanthramides, Novel Substituted N-Cinnamoylanthranilate Alkaloids from Oat Groats and Hulls. *Journal of Agricultural and Food Chemistry*, 37(1), 60-66.
- Collins, F. W. (1986). Oat phenolics: structure, occurrence, and function. In). Minnesota, US: American Association of Cereal Chemists.
- Collins, F. W., Mclachlan, D. C., & Blackwell, B. A. (1991). Oat Phenolics - Avenalumic Acids, a New Group of Bound Phenolic-Acids from Oat Groats and Hulls. *Cereal Chemistry*, 68(2), 184-189.
- Dar, M. Z., Deepika, K., Jan, K., Swer, T. L., Kumar, P., Verma, R., Verma, K., Prakash, K. S., Jan, S., & Bashir, K. (2018). Modification of structure and physicochemical properties of buckwheat and oat starch by gamma-irradiation. *International Journal of Biological Macromolecules*, 108, 1348-1356.
- Dawkins, N. L., & Nnanna, I. A. (1995). Studies on Oat Gum [(1-3,1-4)-Beta-D-Glucan] -Composition, Molecular-Weight Estimation and Rheological Properties. *Food Hydrocolloids*, 9(1), 1-7.
- de Bruijn, W. J. C., van Dinteren, S., Gruppen, H., & Vincken, J. P. (2019). Mass spectrometric characterisation of avenanthramides and enhancing their production by germination of oat (*Avena sativa*). *Food Chemistry*, 277, 682-690.
- Dietrych-Szostak, D., & Oleszek, W. (1999). Effect of processing on the flavonoid content in buckwheat (*Fagopyrum esculentum* Moench) grain. *Journal of Agricultural and Food Chemistry*, 47(10), 4384-4387.
- Dimberg, L. H., Gissen, C., & Nilsson, J. (2005). Phenolic compounds in oat grains (*Avena sativa* L.) grown in conventional and organic systems. *AMBIO: A Journal of the Human Environment*, 34(4-5), 331-337.
- Dimburg, L. H., Theander, O., & Lingnert, H. (1993). Avenanthramides-A group of phenolic antioxidants in oats. *Cereal Chemistry*, 70, 637-641.
- Doehlert, D. C., Simsek, S., Thavarajah, D., Thavarajah, P., & Ohm, J. B. (2013). Detailed Composition Analyses of Diverse Oat Genotype Kernels Grown in Different Environments in North Dakota. *Cereal Chemistry*, 90(6), 572-578.
- Dokuyucu, T., Peterson, D. M., & Akkaya, A. (2003). Contents of antioxidant compounds in Turkish oats: Simple phenolics and avenanthramide concentrations. *Cereal Chemistry*, 80(5), 542-543.
- Dorrell, D. G. (1971a). Fatty acid composition of buckwheat seed. *Journal of the American Oil Chemists' Society*, 48(11), 693 -696
- Dorrell, D. G. (1971b). Fatty Acid Composition of Buckwheat Seed. *Journal of the American Oil Chemists Society*, 48(11), 693-&.
- Duke, J. A. (1992). *Handbook of Phytochemicals Constituents of GRAS Herbs and Other Economic Plants*. Boca Raton, FL: CRC Press,.
- Duve, K. J., & White, P. J. (1991). Extraction and Identification of Antioxidants in Oats. *Journal of the American Oil Chemists Society*, 68(6), 365-370.
- Dziadek, K., Kopec, A., Pastucha, E., Piatkowska, E., Leszczynska, T., Pisulewska, E., Witkowicz, R., & Francik, R. (2016). Basic chemical composition and bioactive compounds content in selected cultivars of buckwheat whole seeds, dehulled seeds and hulls. *Journal of Cereal Science*, 69, 1-8.
- Dziedzic, K., Gorecka, D., Szwengiel, A., Sulewska, H., Kreft, I., Gujska, E., & Walkowiak, J. (2018). The Content of Dietary Fibre and Polyphenols in Morphological Parts of Buckwheat (*Fagopyrum tataricum*). *Plant Foods for Human Nutrition*, 73(1), 82-88.
- Eichenberger Wa U, B. (1984). Sterols in seeds and leaves of oats (*Avena sativa* L.). *Plant Cell Reports*, 3(6), 226-229.
- Emmons, C. L., & Peterson, D. M. (1999). Antioxidant activity and phenolic contents of oat groats and hulls. *Cereal Chemistry*, 76(6), 902-906.

- Fabjan, N., Rode, J., Kosir, I. J., Wang, Z., Zhang, Z., & Kreft, I. (2003). Tartary buckwheat (*Fagopyrum tataricum* Gaertn.) as a source of dietary rutin and quercitrin. *Journal of Agricultural and Food Chemistry*, 51(22), 6452-6455.
- Flander L, S.-M. M., Suortti T, Autio K (2007). Optimization of ingredients and baking process for improved wholemeal oat bread quality. *LWT- Food Science and Technology*, 40, 860–870.
- Food and Agriculture Organization. (2003). Global Information and Early Warning System on Food and Agriculture. In). Rome: FAO.
- Gabrovská, D., Fiedlerová, V., Holasová, M., Mašková, E., Smrčinov, H., Rysová, J., Winterová, R., Michalová, A., & and Hutař, M. (2002). The nutritional evaluation of underutilized cereals and buckwheat. *Food Nutrition Bulletin*, 23, 246-249.
- Gallagher, R. S., Ananth, R., Granger, K., Bradley, B., Anderson, J. V., & Fuerst, E. P. (2010). Phenolic and short-chained aliphatic organic acid constituents of wild oat (*Avena fatua* L.) seeds. *Journal of Agricultural and Food Chemistry*, 58(1), 218-225.
- Gangopadhyay, N., Hossain, M. B., Rai, D. K., & Brunton, N. P. (2015). A Review of Extraction and Analysis of Bioactives in Oat and Barley and Scope for Use of Novel Food Processing Technologies. *Molecules*, 20(6), 10884-10909.
- Garleb, K. A., Bourquin, L. D., Hsu, J. T., Wagner, G. W., Schmidt, S. J., & Fahey, G. C., Jr. (1991). Isolation and chemical analyses of nonfermented fiber fractions of oat hulls and cottonseed hulls. *Journal of Animal Science*, 69(3), 1255-1271.
- Ge, R. H., & Wang, H. (2020). Nutrient components and bioactive compounds in tartary buckwheat bran and flour as affected by thermal processing. *International Journal of Food Properties*, 23(1), 127-137.
- Golijan, J., Milincic, D. D., Petronijevic, R., Pesic, M. B., Barac, M. B., Secanski, M., Lekic, S., & Kostic, A. Z. (2019). The fatty acid and triacylglycerol profiles of conventionally and organically produced grains of maize, spelt and buckwheat. *Journal of Cereal Science*, 90, 102845.
- Grundy, M. M., Fardet, A., Tosh, S. M., Rich, G. T., & Wilde, P. J. (2018). Processing of oat: the impact on oat's cholesterol lowering effect. *Food & Function*, 9(3), 1328-1343.
- Gulpinar, A. R., Orhan, I. E., Kan, A., Senol, F. S., Celik, S. A., & Kartal, M. (2011). Estimation of invitro neuroprotective properties and quantification of rutin and fatty acids in buckwheat (*Fagopyrum esculentum* Moench) cultivated in Turkey. *Food Research International*, 46(2), 536-543.
- Gulpinar, A. R., Orhan, I. E., Kan, A., Senol, F. S., Celik, S. A., & Kartal, M. (2011). Estimation of invitro neuroprotective properties and quantification of rutin and fatty acids in buckwheat (*Fagopyrum esculentum* Moench) cultivated in Turkey. . *Food Research International*, 46(2), 536-543.
- Gunther-Jordanland, K., Dawid, C., Dietz, M., & Hofmann, T. (2016). Key Phytochemicals Contributing to the Bitter Off-Taste of Oat (*Avena sativa* L.). *Journal of Agricultural and Food Chemistry*, 64(51), 9639-9652.
- Guo, X., Zhu, K., Zhang, H., & Yao, H. (2007). Purification and characterization of the antitumor protein from Chinese tartary buckwheat (*Fagopyrum tataricum* Gaertn.) water-soluble extracts. *Journal of Agricultural and Food Chemistry*, 55(17), 6958-6961.
- Guo, X. D., Ma, Y. J., Parry, J., Gao, J. M., Yu, L. L., & Wang, M. (2011). Phenolics content and antioxidant activity of tartary buckwheat from different locations. *Molecules*, 16(12), 9850-9867.
- Habtemariam, S. (2019). Antioxidant and Rutin Content Analysis of Leaves of the Common Buckwheat (*Fagopyrum esculentum* Moench) Grown in the United Kingdom: A Case Study. *Antioxidants (Basel)*, 8(6), 10.
- Hamberg, M., Liepinsh, E., Otting, G., & Griffiths, W. (1998). Isolation and structure of a new galactolipid from oat seeds. *Lipids*, 33(4), 355-363.

- Hitayezu, R., Baakdah, M. M., Kinnin, J., Henderson, K., & Tsopmo, A. (2015). Antioxidant activity, avenanthramide and phenolic acid contents of oat milling fractions. *Journal of Cereal Science*, 63, 35-40.
- Holasova, M., Fiedlerova, V., Smrcinova, H., Orsak, M., Lachman, J., & Vavreinova, S. (2002). Buckwheat the source of antioxidant activity in functional foods. *Food Research International*, 35(2-3), 207-211.
- Holasovaa, M., Fiedlerovaa, V., Smrcinovaa, H., Orsakb, M., Lachmanb, J., & Vavreinovaa, S. (2002). Buckwheat-the source of antioxidant activity in functional foods. *Food Research International*, 35(2-3), 207-211.
- Holmback, J., Karlsson, A. A., & Arnoldsson, K. C. (2001). Characterization of N-acylphosphatidylethanolamine and acylphosphatidylglycerol in oats. *Lipids*, 36(2), 153-165.
- Horbowicz, M., & Obendorf, R. L. (1992). Changes in Sterols and Fatty-Acids of Buckwheat Endosperm and Embryo during Seed Development. *Journal of Agricultural and Food Chemistry*, 40(5), 745-750.
- Hozová, B., Kuniak, L., Moravčíková, P., & Gajdošová, A. (2008). Determination of water-insoluble &beta;-D-glucan in the whole-grain cereals and pseudocereals. *Czech Journal of Food Sciences*, 25(No. 6), 316-324.
- Hu, C., Tang, Y., Zhao, Y., & Sang, S. (2019). Quantitative Analysis and Anti-inflammatory Activity Evaluation of the A-Type Avenanthramides in Commercial Sprouted Oat Products. *Journal of Agricultural and Food Chemistry*.
- Hung, P. V., & Morita, N. (2008). Distribution of phenolic compounds in the graded flours milled from whole buckwheat grains and their antioxidant capacities. *Food Chemistry*, 109(2), 325-331.
- Ikeda, S., Tomura, K., & Kreft, I. (2002). Nutritional characteristics of iron in buckwheat flour. *Fagopyrum*, 19, 79-82.
- Ikeda, S., Yamashita, Y., Kusumoto, K., & Kreft, I. (2005). Nutritional characteristics of minerals in various buckwheat groats. *Fagopyrum*, 22, 71-75.
- Ikeda, S., Yamashita, Y., Tomura, K., & Kreft, I. (2006). Nutritional comparison in mineral characteristics between buckwheat and cereals. *Fagopyrum*, 23, 61-65.
- Ishihara, A., Kojima, K., Fujita, T., Yamamoto, Y., & Nakajima, H. (2014). New series of avenanthramides in oat seed. *Bioscience, Biotechnology, and Biochemistry*, 78(12), 1975-1983.
- Jiang, S., Liu, Q., Xie, Y., Zeng, H., Zhang, L., Jiang, X., & Chen, X. (2015). Separation of five flavonoids from tartary buckwheat (*Fagopyrum tataricum* (L.) Gaertn) grains via off-line two dimensional high-speed counter-current chromatography. *Food Chemistry*, 186, 153-159.
- Kalinova, J., Triska, J., & Vrchotova, N. (2006). Distribution of Vitamin E, squalene, epicatechin, and rutin in common buckwheat plants (*Fagopyrum esculentum* Moench). *Journal of Agricultural and Food Chemistry*, 54(15), 5330-5335.
- Keriene, I., Mankeviciene, A., Bliznikas, S., Jablonskyte-Rasce, D., Maiksteniene, S., & Cesnuleviciene, R. (2015). Biologically active phenolic compounds in buckwheat, oats and winter spelt wheat. *Zemdirbyste-Agriculture*, 102(3), 289-296.
- Khalass, S., Popescu, C., Popescu, I. A., Arsene, A. L., Udeanu, D. I., & Lupuliasa, D. (2018). Physico-Chemical and Microbiological Characterization of *Fagopyrum Esculentum* Moench. Fatty Oil Obtained by Cold Pressure. *Farmacia*, 66(5), 899-903.
- Kim, H. J., Park, K. J., & Lim, J. H. (2011). Metabolomic analysis of phenolic compounds in buckwheat (*Fagopyrum esculentum* M.) sprouts treated with methyl jasmonate. *Journal of Agricultural and Food Chemistry*, 59(10), 5707-5713.
- Kim, S. J., Zaidul, I. S., Suzuki, T., Mukasa, Y., Hashimoto, N., Takigawa, S., Noda, T., Matsuura-Endo, C., & Yamauchi, H. (2008). Comparison of phenolic compositions between common and tartary buckwheat (*Fagopyrum*) sprouts. *Food Chemistry*, 110(4), 814-820.



- Kim, S. L., Kim, S. K., and Park, C. H. (2002). Comparisons of lipid, fatty acids and tocopherols of different buckwheat species. *Food Science and Biotechnology*, 11, 322-336.
- Klose, C., Schehl, B. D., & Arendt, E. K. (2009). Fundamental study on protein changes taking place during malting of oats. *Journal of Cereal Science*, 49(1), 83-91.
- Kočevár Glavač, N., Stojilkovski, K., Kreft, S., Park, C. H., & Kreft, I. (2017). Determination of fagopyrins, rutin, and quercetin in Tartary buckwheat products. *LWT - Food Science and Technology*, 79, 423-427.
- Kovacova, M., & Malinova, E. (2007). Ferulic and coumaric acids, total phenolic compounds and their correlation in selected oat genotypes. *Czech Journal of Food Sciences*, 25(6), 325-332.
- Kraujaliene, V., Pukalskas, A., & Venskutonis, P. R. (2017). Multi-stage recovery of phytochemicals from buckwheat (*Fagopyrum esculentum* Moench) flowers by supercritical fluid and pressurized liquid extraction methods. *Industrial Crops and Products*, 107, 271-280.
- Kreft, S., Janes, D., & Kreft, I. (2013). The content of fagopyrin and polyphenols in common and tartary buckwheat sprouts. *Acta Pharmaceutica*, 63(4), 553-560.
- Kreft, S., Knapp, M., & Kreft, I. (1999). Extraction of rutin from buckwheat (*Fagopyrum esculentum* Moench) seeds and determination by capillary electrophoresis. *Journal of Agricultural and Food Chemistry*, 47(11), 4649-4652.
- Krskoskova, B., & Mrazova, Z. (2005). Prophylactic components of buckwheat. *Food Research International*, 38(5), 561-568.
- Krumina-Zemture, G., & Beitane, I. (2018). *Fatty acid composition in buckwheat (fagopyrum esculentum m.) Flours and their extruded products*. Kaunas, Lithuania.
- Larsson, M., & Sandberg, A. S. (1992). Phytate Reduction in Oats during Malting. *Journal of Food Science*, 57(4), 994-997.
- Lee, J. M., Lee, K. H., Yoon, Y. H., Cho, E. J., & Lee, S. (2013). Identification of triterpenoids and flavonoids from the seeds of tartary buckwheat. *Natural Product Sciences*, 19(2), 137-144.
- Lee, K. Y., Park, K. H., Ahn, J. B., & Lee, H. G. (2014). Effect of different chemical modification on the physicochemical properties of fiber-enriched polysaccharides isolated from wholegrain rice and buckwheat. *Food Science and Biotechnology*, 23(5), 1469-1475.
- Leonova, S., Shelenga, T., Hamberg, M., Konarev, A. V., Loskutov, I., & Carlsson, A. S. (2008). Analysis of oil composition in cultivars and wild species of oat (*Avena* sp.). *Journal of Agricultural and Food Chemistry*, 56(17), 7983-7991.
- Li, C. H., Matsui, T., Matsumoto, K., Yamasaki, R., & Kawasaki, T. (2002). Latent production of angiotensin I-converting enzyme inhibitors from buckwheat protein. *Journal of Peptide Science*, 8(6), 267-274.
- Li, F., Yuan, Y., Yang, X., Tao, S., & Ming, J. (2013). Phenolic Profiles and Antioxidant Activity of Buckwheat (*Fagopyrum esculentum* Möench and *Fagopyrum tartaricum* L. Gaerth) Hulls, Brans and Flours. *Journal of Integrative Agriculture*, 12(9), 1684-1693.
- Li, L., Lietz, G., & Seal, C. (2018). Buckwheat and CVD Risk Markers: A Systematic Review and Meta-Analysis. *Nutrients*, 10(5).
- Lin, H. C., & Lin, J. Y. (2016). Immune Cell-Conditioned Media Suppress Prostate Cancer PC-3 Cell Growth Correlating With Decreased Proinflammatory/Anti-inflammatory Cytokine Ratios in the Media Using 5 Selected Crude Polysaccharides. *Integrative Cancer Therapies*, 15(4), NP13-NP25.
- Lintschinger, J., Fuchs, N., Moser, H., Jager, R., Hlebeina, T., Markolin, G., & Gossler, W. (1997). Uptake of various trace elements during germination of wheat, buckwheat and quinoa. *Plant Foods for Human Nutrition*, 50(3), 223-237.
- Liu, G., Li, J., Shi, K., Wang, S., Chen, J., Liu, Y., & Huang, Q. (2009). Composition, secondary structure, and self-assembly of oat protein isolate. *Journal of Agricultural and Food Chemistry*, 57(11), 4552-4558.

- Liu, Y., Cai, C., Yao, Y., & Xu, B. (2019). Alteration of phenolic profiles and antioxidant capacities of common buckwheat and tartary buckwheat produced in China upon thermal processing. *Journal of the science of food and agriculture*, 99(12), 5565-5576.
- Maatta, K., Lampi, A. M., Petterson, J., Fogelfors, B. M., Piironen, V., & Kamal-Eldin, A. (1999). Phytosterol content in seven oat cultivars grown at three locations in Sweden. *Journal of the science of food and agriculture*, 79(7), 1021-1027.
- Manthey, F. A., Hareland, G. A., & Huseby, D. J. (1999). Soluble and insoluble dietary fiber content and composition in oat. *Cereal Chemistry*, 76(3), 417-420.
- Mary, W., Crombie, L., & Crombie, L. (1986). Distribution of avenacins A-1, A-2, B-1 and B-2 in oat roots: Their fungicidal activity towards 'take-all' fungus. *Phytochemistry*, 25(9), 2069-2073.
- Mattila, P., Pihlava, J. M., & Hellstrom, J. (2005). Contents of phenolic acids, alkyl- and alkenylresorcinols, and avenanthramides in commercial grain products. *Journal of Agricultural and Food Chemistry*, 53(21), 8290-8295.
- Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., & Group, P. (2009). Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*, 6(7), e1000097.
- Moreau, R. A., Doehlert, D. C., Welti, R., Isaac, G., Roth, M., Tamura, P., & Nuñez, A. (2008). The Identification of Mono-, Di-, Tri-, and Tetragalactosyldiacylglycerols and their Natural Estolides in Oat Kernels. *Lipids*, 43(6), 533-548.
- Muka, T., Glisic, M., Milic, J., Verhoog, S., Bohlius, J., Bramer, W., Chowdhury, R., & Franco, O. H. (2020). A 24-step guide on how to design, conduct, and successfully publish a systematic review and meta-analysis in medical research. *European Journal of Epidemiology*, 35(1), 49-60.
- Multari, S., Pihlava, J. M., Ollennu-Chuasam, P., Hietaniemi, V., Yang, B., & Suomela, J. P. (2018). Identification and Quantification of Avenanthramides and Free and Bound Phenolic Acids in Eight Cultivars of Husked Oat ( *Avena sativa* L) from Finland. *Journal of Agricultural and Food Chemistry*, 66(11), 2900-2908.
- Musa Ozcan, M., Ozkan, G., & Topal, A. (2006). Characteristics of grains and oils of four different oats (*Avena sativa* L.) cultivars growing in Turkey. *International Journal of Food Sciences and Nutrition*, 57(5-6), 345-352.
- Nam, T. G., Lee, S. M., Park, J. H., Kim, D. O., Baek, N. I., & Eom, S. H. (2015). Flavonoid analysis of buckwheat sprouts. *Food Chemistry*, 170, 97-101.
- Obendorf, R. L. (1997). Oligosaccharides and galactosyl cyclitols in seed desiccation tolerance. *Seed Science Research*, 7(2), 63-74.
- Obendorf, R. L., Steadman, K. J., Fuller, D. J., Horbowicz, M., & Lewis, B. A. (2000). Molecular structure of fagopyritol A1 (O- $\alpha$ -d-galactopyranosyl-(1 $\rightarrow$ 3)-d-chiro-inositol) by NMR. *Carbohydrate Research*, 328(4), 623-627.
- Onning, G., & Asp, N. G. (1993). Analysis of Saponins in Oat Kernels. *Food Chemistry*, 48(3), 301-305.
- Oomah, B. D., & Mazza, G. (1996). Flavonoids and antioxidative activities in buckwheat. *Journal of Agricultural and Food Chemistry*, 44(7), 1746-1750.
- Oomah, B. D., & Mazza, G. (1996). Flavonoids and Antioxidative Activities in Buckwheat. *Journal of Agricultural and Food Chemistry*, 44(7), 1746-1750.
- Ortiz-Robledo, F., Villanueva-Fierro, I., Oomah, B. D., Lares-Asef, I., Proal-Najera, J. B., & Navar-Chaidez, J. J. (2013). Avenanthramides and nutritional components of four Mexican oat (*Avena sativa* L.) varieties. *Agrociencia*, 47(3), 225-232.
- Osborn, A. E., Clarke, B. R., Lunness, P., Scott, P. R., & Daniels, M. J. (1994). An Oat Species Lacking Avenacin Is Susceptible to Infection by *Gaeumannomyces Graminis* Var *Tritici*. *Physiological and Molecular Plant Pathology*, 45(6), 457-467.
- Panfili, G., Fratianni, A., Criscio, T. D., & Marconi, E. (2008). Tocol and  $\beta$ -glucan levels in barley varieties and in pearling by-products. *Food Chemistry*(107), 84-91.

612 Pankaja, S. (2011). Phenolic contents, antioxidant and  $\alpha$ -glucosidase inhibition properties of Nepalese  
613 strain buckwheat vegetables. *African Journal of Biotechnology*, 11(1).

614 Park, C. H., Yeo, H. J., Park, Y. E., Chun, S. W., Chung, Y. S., Lee, S. Y., & Park, S. U. (2019). Influence of  
615 Chitosan, Salicylic Acid and Jasmonic Acid on Phenylpropanoid Accumulation in Germinated 616  
Buckwheat (*Fagopyrum esculentum* Moench). *Foods*, 8(5), 10.

617 Pecio, L., Wawrzyniak-Szolowska, A., Oleszek, W., & Stochmal, A. (2013). Rapid analysis of  
618 avenacosides in grain and husks of oats by UPLC-TQ-MS. *Food Chemistry*, 141(3), 2300-2304.

619 Peng, L., Zou, L., Tan, M., Deng, Y., Yan, J., Yan, Z., & Zhao, G. (2017). Free amino acids, fatty acids and  
620 phenolic compounds in Tartary buckwheat of different hull colour. *Czech Journal of Food*  
621 *Sciences*, 35(3), 214-222.

622 Peterson, D. M., & Dimberg, L. H. (2008). Avenanthramide concentrations and hydroxycinnamoyl-  
623 CoA : hydroxyanthranilate N-hydroxycinnamoyltransferase activities in developing oats.  
624 *Journal of Cereal Science*, 47(1), 101-108.

625 Peterson, D. M., Jensen, C. M., Hoffman, D. L., & Mannerstedt-Fogelfors, B. (2007). Oat tocots:  
626 Saponification vs. direct extraction and analysis in high-oil genotypes. *Cereal Chemistry*,  
627 84(1), 56-60.

628 Peterson, D. M., & Qureshi, A. A. (1993). Genotype and environment effects on tocots of barley and  
629 oats. *Cereal Chemistry*, 70, 157-162.

630 Pridal, A. A., Bottger, W., & Ross, A. B. (2018). Analysis of avenanthramides in oat products and  
631 estimation of avenanthramide intake in humans. *Food Chemistry*, 253, 93-100.

632 Qin, P. Y., Wang, Q., Shan, F., Hou, Z. H., & Ren, G. X. (2010). Nutritional composition and flavonoids  
633 content of flour from different buckwheat cultivars. *International Journal of Food Science and* 634  
*Technology*, 45(5), 951-958.

635 Quettier-Deleu, C., Gressier, B., Vasseur, J., Dine, T., Brunet, C., Luyckx, M., Cazin, M., Cazin, J.-C.,  
636 Bailleul, F., & Trotin, F. (2000). Phenolic compounds and antioxidant activities of buckwheat  
637 (*Fagopyrum esculentum* Moench) hulls and flour. *Journal of ethnopharmacology*, 72(1-2), 35-  
638 42.

639 Rasane, P., Jha, A., Sabikhi, L., Kumar, A., & Unnikrishnan, V. S. (2015). Nutritional advantages of oats  
640 and opportunities for its processing as value added foods - a review. *Journal of Food Science*  
641 *and Technology*, 52(2), 662-675.

642 Regand, A., Chowdhury, Z., Tosh, S. M., Wolever, T. M. S., & Wood, P. (2011). The molecular weight,  
643 solubility and viscosity of oat beta-glucan affect human glycemic response by modifying  
644 starch digestibility. *Food Chemistry*, 129(2), 297-304.

645 Ren, Q., Wu, C., Ren, Y., & Zhang, J. (2013). Characterization and identification of the chemical  
646 constituents from tartary buckwheat (*Fagopyrum tataricum* Gaertn) by high performance  
647 liquid chromatography/photodiode array detector/linear ion trap FTICR hybrid mass  
648 spectrometry. *Food Chemistry*, 136(3-4), 1377-1389.

649 Rice-Evans, C., & Paganga, G. (1997). Antioxidant properties of phenolic compounds. *Trends Plant*  
650 *Science*, 2, 152-159.

651 Runyon, J. R., Nilsson, L., Alftren, J., & Bergenstahl, B. (2013). Characterization of oat proteins and  
652 aggregates using asymmetric flow field-flow fractionation. *Analytical and Bioanalytical*  
653 *Chemistry*, 405(21), 6649-6655.

654 Sadauskiene, I., Liekis, A., Bernotiene, R., Sulinskiene, J., Kasauskas, A., & Zekonis, G. (2018). The  
655 Effects of Buckwheat Leaf and Flower Extracts on Antioxidant Status in Mouse Organs.  
656 *Oxidative Medicine and Cellular Longevity*, 2018, 6712407.

657 Sang, S., & Chu, Y. (2017). Whole grain oats, more than just a fiber: Role of unique phytochemicals.  
658 *Mol Nutr Food Res*, 61(7).

659 Saxena, V. K. S., G.C. (1987). A new flavonoid from *Fagopyrum tataricum*. *Fitoterapia*, 58283.

- 660 Sedej, I., Sakac, M., Mandic, A., Misan, A., Tumbas, V., & Canadanovic-Brunet, J. (2012). Buckwheat  
661 (*Fagopyrum esculentum* Moench) grain and fractions: antioxidant compounds and activities.  
662 *Journal of Food Science*, 77(9), C954-959.
- 663 Shao, M., Yang, Y. H., Gao, H. Y., Wu, B., Wang, L. B., & Wu, L. J. (2005). Studies on the chemical  
664 constituents of *Fagopyrum dibotrys*. *Journal of Shenyang Pharmaceutical University*, 22, 100-  
665 102.
- 666 Shewry, P. R., Piironen, V., Lampi, A. M., Nystrom, L., Li, L., Rakszegi, M., Fras, A., Boros, D., Gebruers,  
667 K., Courtin, C. M., Delcour, J. A., Andersson, A. A., Dimberg, L., Bedo, Z., & Ward, J. L. (2008).  
668 Phytochemical and fiber components in oat varieties in the HEALTHGRAIN Diversity Screen.  
669 *Journal of Agricultural and Food Chemistry*, 56(21), 9777-9784.
- 670 Shi, J., Arunasalam, K., Yeung, D., Kakuda, Y., Mittal, G., & Jiang, Y. (2004). Saponins from edible  
671 legumes: chemistry, processing, and health benefits. *Journal of medicinal food*, 7(1), 67-78.
- 672 Sinkovic, L., Kokalj, D., Vidrih, R., & Meglic, V. (2020a). Milling fractions fatty acid composition of  
673 common (*Fagopyrum esculentum* Moench) and tartary (*Fagopyrum tataricum* (L.) Gaertn)  
674 buckwheat. *Journal of Stored Products Research*, 85, 5.
- 675 Sinkovic, L., Kokalj, D., Vidrih, R., & Meglic, V. (2020b). Milling fractions fatty acid composition of  
676 common (*Fagopyrum esculentum* Moench) and tartary (*Fagopyrum tataricum* (L.) Gaertn)  
677 buckwheat. *Journal of Stored Products Research*, 85, 101551.
- 678 Siwatch, M., Yadav, R. B., & Yadav, B. S. (2019). Influence of Processing Treatments on Nutritional  
679 and Physicochemical Characteristics of Buckwheat (*Fagopyrum esculentum*). *Current*  
680 *Nutrition & Food Science*, 15(4), 408-414.
- 681 Skoglund, M., Peterson, D. M., Andersson, R., Nilsson, J., & Dimberg, L. H. (2008). Avenanthramide  
682 content and related enzyme activities in oats as affected by steeping and germination.  
683 *Journal of Cereal Science*, 48(2), 294-303.
- 684 Smeds, A. I., Jauhiainen, L., Tuomola, E., & Peltonen-Sainio, P. (2009). Characterization of variation in  
685 the lignan content and composition of winter rye, spring wheat, and spring oat. *Journal of*  
686 *Agricultural and Food Chemistry*, 57(13), 5837-5842.
- 687 Sosulski, F., Krygier, K., & Hogge, L. (1982). Free, Esterified, and Insoluble-Bound Phenolic-Acids .3.  
688 Composition of Phenolic-Acids in Cereal and Potato Flours. *Journal of Agricultural and Food*  
689 *Chemistry*, 30(2), 337-340.
- 690 Sowa, S. M. H., & White, P. J. (1992). Characterization of Starch Isolated from Oat Groats with  
691 Different Amounts of Lipid. *Cereal Chemistry*, 69(5), 521-527.
- 692 Soycan, G., Schar, M. Y., Kristek, A., Boberska, J., Alsharif, S. N. S., Corona, G., Shewry, P. R., &  
693 Spencer, J. P. E. (2019). Composition and content of phenolic acids and avenanthramides in  
694 commercial oat products: Are oats an important polyphenol source for consumers? *Food*  
695 *Chemistry*, 3, 100047.
- 696 Steadman, K. J., Burgoon, M. S., Lewis, B. A., Edwardson, S. E., & Obendorf, R. L. (2001). Minerals,  
697 phytic acid, tannin and rutin in buckwheat seed milling fractions. *Journal of the science of*  
698 *food and agriculture*, 81(11), 1094-1100.
- 699 Steadman, K. J., Burgoon, M. S., Schuster, R. L., Lewis, B. A., Edwardson, S. E., & Obendorf, R. L.  
700 (2000). Fagopyritols, D-chiro-inositol, and other soluble carbohydrates in buckwheat seed  
701 milling fractions. *Journal of Agricultural and Food Chemistry*, 48(7), 2843-2847.
- 702 Steadman, K. J., Fuller, D. J., & Obendorf, R. L. (2001). Purification and molecular structure of two  
703 digalactosyl D-chiro-inositols and two trigalactosyl D-chiro-inositols from buckwheat seeds.  
704 *Carbohydrate Research*, 331, 19-25.
- 705 Steer, M. W., Gunning, B. E., Graham, T. A., & Carr, D. J. (1968). Isolation, properties, and structure of  
706 fraction I protein from *Avena sativa* L. *Planta*, 79(3), 254-267.
- 707 Sytar, O. (2015). Phenolic acids in the inflorescences of different varieties of buckwheat and their  
708 antioxidant activity. *Journal of King Saud University - Science*, 27(2), 136-142.

709 Takahama, U., & Hirota, S. (2010). Fatty acids, epicatechin-dimethylgallate, and rutin interact with  
710 buckwheat starch inhibiting its digestion by amylase: implications for the decrease in 711  
glycemic index by buckwheat flour. *Journal of Agricultural and Food Chemistry*, 58(23), 712  
12431-12439.

713 Tanwar, B., & Modgil, R. (2012). Flavonoids: Dietary Occurrence and Health Benefits. *Spatula DD -*  
714 *Peer Reviewed Journal on Complementary Medicine and Drug Discovery*, 2(1), 59-68.

715 Terpinc, P., Cigic, B., Polak, T., Hribar, J., & Pozrl, T. (2016). LC-MS analysis of phenolic compounds  
716 and antioxidant activity of buckwheat at different stages of malting. *Food Chemistry*, 210, 9-  
717 17.

718 Tian, L., & Xu, L. Z. (1997). Studies on the chemical constituents of the aerial parts of *Fagopyrum*  
719 *dibotrys*. *China Journal of Chinese Materia Medica*, 22, 743-745.

720 Tien, N. N. T., Trinh, L. N. D., Inoue, N., Morita, N., & Hung, P. V. (2018). Nutritional composition,  
721 bioactive compounds, and diabetic enzyme inhibition capacity of three varieties of  
722 buckwheat in Japan. *Cereal Chemistry*, 95(5), 615-624.

723 Tong, L. T., Liu, L. Y., Zhong, K., Wang, Y., Guo, L. N., & Zhou, S. M. (2014). Effects of Cultivar on  
724 Phenolic Content and Antioxidant Activity of Naked Oat in China. *Journal of Integrative* 725  
*Agriculture*, 13(8), 1809-1816.

726 Trojanowska, M. R., Osbourn, A. E., Daniels, M. J., & Threlfall, D. R. (2000). Biosynthesis of avenacins  
727 and phytosterols in roots of *Avena sativa* cv. Image. *Phytochemistry*, 54(2), 153-164.

728 Tschesche, R., & Lauven, P. (1971). Steroidsaponine mit mehr als einer Zuckerkette, V. Avenacosid B,  
729 ein zweites bisdesmosidisches Steroidsaponin aus *Avena sativa*. *Chemische Berichte*, 104(11),  
730 3549-3555.

731 Tschesche, R., Tauscher, M., Fehlhaber, H.-W., & Wulff, G. (1969). Steroidsaponine mit mehr als einer  
732 Zuckerkette, IV. Avenacosid A, ein bisdesmosidisches Steroidsaponin aus *Avena sativa*.  
733 *Chemische Berichte*, 102(6), 2072-2082.

734 Tschesche, R., & Wiemann, W. (1977). Steroidsaponine mit mehr als einer Zuckerkette, XI. Desgluco-  
735 avenacosid-A und -B, biologisch aktive Nuatigeninglycoside. *Chemische Berichte*, 110(7),  
736 2416-2423.

737 Tschesche, R., & Wulff, G. (1973). Chemie und Biologie der Saponine. In *Fortschritte Der Chemie*  
738 *Organischer Naturstoffe*, vol. 30 (pp. 461-606).

739 Tsuzuki, W., Ogata, Y., Akasaka, K., Shibata, S., & Suzuki, T. (1991). Fatty-acid composition of selected  
740 buckwheat species by fluorometric high-performance liquid chromatography. *Cereal*  
741 *Chemistry*, 68, 365-369.

742 van den Broeck, H. C., Londono, D. M., Timmer, R., Smulders, M. J., Gilissen, L. J., & van der Meer, I.  
743 M. (2015). Profiling of Nutritional and Health-Related Compounds in Oat Varieties. *Foods*,  
744 5(1), 2.

745 Varga, M., Jojart, R., Fonad, P., Mihaly, R., & Palagyi, A. (2018). Phenolic composition and antioxidant  
746 activity of colored oats. *Food Chemistry*, 268, 153-161.

747 Verardo, V., Arráez-Román, D., Segura-Carretero, A., Marconi, E., Fernández-Gutiérrez, A., & Caboni,  
748 M. F. (2010). Identification of buckwheat phenolic compounds by reverse phase high  
749 performance liquid chromatography–electrospray ionization–time of flight–mass  
750 spectrometry (RP-HPLC–ESI-TOF-MS). *Journal of Cereal Science*, 52(2), 170-176.

751 Verardo, V., Serea, C., Segal, R., & Caboni, M. F. (2011). Free and bound minor polar compounds in  
752 oats: Different extraction methods and analytical determinations. *Journal of Cereal Science*,  
753 54(2), 211-217.

754 Wang, K. J., Zhang, Y. J., & Yang, C. R. (2005). Antioxidant phenolic constituents from *Fagopyrum*  
755 *dibotrys*. *Journal of ethnopharmacology*, 99(2), 259-264.

756 Wang, X. T., Zhu, Z. Y., Zhao, L., Sun, H. Q., Meng, M., Zhang, J. Y., & Zhang, Y. M. (2016). Structural  
757 characterization and inhibition on alpha-d-glucosidase activity of non-starch polysaccharides  
758 from *Fagopyrum tartaricum*. *Carbohydrate Polymers*, 153, 679-685.

Wang, Z., Wang, L., Chang, W., Li, Y., Zhang, Z., Wieslander, G., & Norback, D. (2006). Cloning, expression, and identification of immunological activity of an allergenic protein in tartary buckwheat. *Bioscience, Biotechnology, and Biochemistry*, 70(5), 1195-1199.

Wang, Z., Zhang, Z., Zhao, Z., Wieslander, G., Norback, D., & Kreft, I. (2004). Purification and characterization of a 24 kDa protein from tartary buckwheat seeds. *Bioscience, Biotechnology, and Biochemistry*, 68(7), 1409-1413.

Watanabe, M., Ohshita, Y., & Tsushida, T. (1997). Antioxidant compounds from buckwheat (*Fagopyrum esculentum* Moench) hulls. *Journal of Agricultural and Food Chemistry*, 45(4), 1039-1044.

Wefers, D., & Bunzel, M. (2015). Characterization of Dietary Fiber Polysaccharides from Dehulled Common Buckwheat (*Fagopyrum esculentum*) Seeds. *Cereal Chemistry*, 92(6), 598-603.

Welch, R. W. (1975). Fatty acid composition of grain from winter and spring sown oats, barley and wheat. *Journal of the Science of Food and Agriculture*, 26(4), 429-435.

Welch, R. W. (1975). Fatty acid composition of grain from winter and spring sown oats, barley and wheat. *Journal of the science of food and agriculture*, 26(4), 429-435.

Wenzig, E., Kunert, O., Ferreira, D., Schmid, M., Schuhly, W., Bauer, R., & Hiermann, A. (2005). Flavonolignans from *Avena sativa*. *Journal of Natural Products*, 68(2), 289-292.

Wu, H. Z., Zhou, J. Y., & Pan, H. L. (2008). Study on chemical constituents of *Fagopyrum dibotrys* (Don) Hara *Chinese Journal of Hospital Pharmacy*, 28, 21-26.

Xie, Z., Mui, T., Sintara, M., Ou, B., Johnson, J., Chu, Y., O'Shea, M., Kasturi, P., & Chen, Y. (2017). Rapid quantitation of avenanthramides in oat-containing products by high-performance liquid chromatography coupled with triple quadrupole mass spectrometry (HPLC-TQMS). *Food Chemistry*, 224, 280-288.

Xing, Y. M., & White, P. J. (1997). Identification and function of antioxidants from oat greats and hulls. *Journal of the American Oil Chemists Society*, 74(3), 303-307.

Xu, B., Xiao, G., & Ding, X. (2002). Determination of phenolic acids and proanthocyanidin in buckwheat (*Fagopyrum tataricum* (Linn) Gaench). *Food and Fermentation Industries*, 28, 32-33.

Yan, X. C., Sun, Z. J., Xie, G., He, Y., Yi, C., & Fang, C. (2011). Purification and monosaccharide composition analysis of tartary buckwheat polysaccharides. *Food Science*, 32, 33-36.

York, W., Darvill, A. G., McNeill, M., & Stevenson, T. (1993). Genotype and environmental effects on tocols of barley and oats. *Cereal Chem*, 70(2), 157-162.

Yu, X., Yang, M., Dong, J., & Shen, R. (2018). Comparative Analysis of the Antioxidant Capacities and Phenolic Compounds of Oat and Buckwheat Vinegars During Production Processes. *J Food Sci*, 83(3), 844-853.

Zemnukhova, L. A., Sukhoverkhov, S. V., Shkorina, E. D., Kovekhova, A. V., & Tomshich, S. V. (2007). Use of high-performance exclusion chromatography for analysis of the molecular mass of polysaccharides produced from buckwheat and sunflower wastes. *Russian Journal of Applied Chemistry*, 80(12), 2051-2056.

Zemnukhova, L. A., Tomshich, S. V., Shkorina, E. D., & Klykov, A. G. (2004). Polysaccharides from buckwheat production wastes. *Russian Journal of Applied Chemistry*, 77(7), 1178-1181.

Zhan-Lu, Z., Mei-Liang, Z., Yu, T., Fa-Liang, L., Yi-Xiong, T., Ji-Rong, S., Wen-Tong, X., & Yan-Min, W. (2012). Bioactive compounds in functional buckwheat food. *Food Research International*, 49, 389-395.

Zhang, C., Lu, Y., Guo, G., & Zhang, H. (2005). Studies on antifatigue of buckwheat protein. *Journal of Food Science and Biotechnology*, 24, 78-82.

Zhang, Q., & Xu, J. G. (2017). Determining the geographical origin of common buckwheat from China by multivariate analysis based on mineral elements, amino acids and vitamins. *Scientific reports*, 7(1), 9696.

808 Zhang, W. K., Xu, J. K., Zhang, L., & Du, G. H. (2012). Flavonoids from the bran of *Avena sativa*.  
809 *Chinese Journal of Natural Medicines*, 10, 110-114.

810 Zhang, Z., Wang, Z., Liu, F., & Lin, R. (1999). Studies on nutrition and anti-senescence function of  
811 protein complex from tartary buckwheat. *Acta Nutrimenta Sinica*, 21, 159-162.

812 Zheng, C., Hu, C., Ma, X., Peng, C., Zhang, H., & Qin, L. (2012). Cytotoxic phenylpropanoid glycosides  
813 from *Fagopyrum tataricum* (L.) Gaertn. *Food Chemistry*, 132(1), 433-438.

814 Zheng, F., Song, Q. B., Qiang, G. R., Sun, P. L., Zhu, X. Y., & Chen, D. X. (2004). GC/MS analysis of fatty  
815 acid in *Fagopyrum esculentum* seed oil. *Food Science*, 25, 267-269.

816

Figure 1. Flowchart of studies included in current review






Figure 2. Illustrative summary of the most important findings

Phytochemical	Oat		Buckwheat		<div><div></div>Oat</div> <div><div></div>Buckwheat</div>
	Total content	Most common compound	Total content	Most common compound	
Avenanthramides	0.5 - 71.85 mg/100g	AVA-A, AVA-B and AVA-C	n.d.	n.a.	<div><div></div></div>
Flavonoids	n.a.	Quercetin (max 8.9 mg/100g); Rutin (max 0.47 mg/100g)	6- 507.6 mg RE/100g	Rutin ( max 5,186 mg/100g); Quercetin (max 857.62 mg/100g)	<div><div></div></div>
Phenolics	180-576 mg RE/100g	Ferulic acids (max 149.36 mg/100g)	123-3,149 mg RE/100g	p-Anisic acid (max 1190 mg/100g)	<div><div></div></div>
Phytosterols	35- 68.2 mg/100g	Sitosterol (59.1-64.9%); campesterol (7.6-9.1%)	19-139 mg/100g	Sitosterol (60-70%); campesterol (5-10%)	<div><div></div></div>
Tocols	0.50-3.61 mg/100g	α-tocotrienol (max 5.64 mg/100g)	5.46-55.22mg/100g	γ-tocopherol (max 22.10 mg/100g)	<div><div></div></div>
β-glucan	5,180- 28,200 mg/100g	n.a	n.a.	1,260-3,500 mg/100g	<div><div></div></div>

The orange and yellow bars present maximal phytochemical content reported in oat vs. maximal phytochemical content reported in buckwheat. For example, for total phenolics, the orange bar illustrates the maximal reported total phenolic content in oat (576 mg RE/100g) and the yellow bar represents the max reported phenolic content in buckwheat (3,149 mg RE/100g).

**Abbreviations:** AVA-A, AVA-B and AVA-C: Esters of 5-hydroxyanthranilic acid with p-coumaric (AVA-A), caffeic (AVA-B) and ferulic acids (AVA-C); n.a.: not available; n.d.: not detected; RE: rutin equivalents

Supplemental table 1. Search strategy used in current review
Embase.com (('metabolism'/exp OR 'metabolomics'/exp OR 'energy metabolism'/exp OR 'metabolic activity assay'/de OR 'phytochemistry'/de OR 'phytochemical'/exp OR 'nutrient content'/de OR 'nutrient'/exp OR 'nutraceutical'/exp OR 'diet supplementation'/de OR 'pharmacokinetics'/exp OR (metabol* OR cometabol* OR co-metabol* OR nutrient* OR nutraceut* OR phytochem* OR phytopharmaceut* OR pharmacokinetic* OR ((dietary OR nutritional) NEAR/3 (supplement*))) :ab,ti) OR ('alkaloid'/exp OR 'amine'/exp OR 'amino acid'/exp OR 'carbohydrate'/exp OR 'carotenoid'/exp OR 'disaccharide'/exp OR 'fatty acid'/exp OR 'flavonoid'/exp OR 'indole derivative'/exp OR 'lignan'/exp OR 'lipid'/exp OR 'monosaccharide'/exp OR 'peptide'/exp OR 'polysaccharide'/exp OR 'protein'/exp OR 'purine derivative'/exp OR 'pyrimidine derivative'/exp OR 'stilbene derivative'/de OR 'sugar alcohol'/exp OR 'terpene'/de OR 'amide'/de OR 'avenanthramide'/exp OR ("Amino acid*" OR "Cyanogenic glucoside*" OR "Fatty acid*" OR "Nucleic acid base*" OR "Organic acid*" OR "Organosulfur compound*" OR "Phenolic acid*" OR "Sugar alcohol*" OR Alkaloid* OR Alkane* OR Amine* OR Benzenoid* OR Carotenoid* OR Chlorophyll* OR Disaccharide* OR Flavonoid* OR Indole* OR Lignan* OR Monosaccharide* OR Peptide* OR Polyacetylene* OR Polysaccharide* OR Protein* OR Purine* OR Pyrimidine* OR Stilbene* OR Terpene* OR Terpenoid* OR avenanthramid* OR amide*) :ab,ti) AND ('plant extract'/de OR 'plant medicinal product'/de OR 'nutrient content'/de OR 'chromatography'/exp OR 'chemistry'/exp OR (chromatogra* OR electrochromatogra* OR ((plant OR plants) NEAR/3 (extract* OR biochem* OR preparation* OR medicinal*))) :ab,ti) AND ('buckwheat'/de OR 'fagopyrum'/exp OR 'oat'/de OR 'oat bran'/de OR (buckwheat* OR fagopyrum OR 'avena sativa' OR 'avena nuda' OR avena OR avenas OR ((oat OR oats) NEAR/3 (plant* OR hull* OR bran* OR kernel* OR groat*))) :ab,ti) NOT ((Conference Abstract]/lim OR [Letter]/lim OR [Note]/lim OR [Editorial]/lim)
Web of Science (Core collection) TS=(((metabol* OR cometabol* OR co-metabol* OR nutrient* OR nutraceut* OR phytochem* OR phytopharmaceut* OR pharmacokinetic* OR ((dietary OR nutritional) NEAR/2 (supplement*)) OR "Amino acid*" OR "Cyanogenic glucosid*" OR "Fatty acid*" OR "Nucleic acid base*" OR "Organic acid*" OR "Organosulfur compound*" OR "Phenolic acid*" OR "Sugar alcohol*" OR Alkaloid* OR Alkane* OR Amine* OR Benzenoid* OR Carotenoid* OR Chlorophyll* OR Disaccharide* OR Flavonoid* OR Indole* OR Lignan* OR Monosaccharide* OR Peptide* OR Polyacetylene* OR Polysaccharide* OR Protein* OR Purine* OR Pyrimidine* OR Stilbene* OR Terpene* OR Terpenoid* OR avenanthramid* OR amide*) AND (chromatogra* OR electrochromatogra* OR ((plant OR plants) NEAR/3 (extract* OR biochem* OR preparation* OR medicinal*)))) AND (buckwheat* OR fagopyrum OR "avena sativa*" OR "avena nuda*" OR avena OR avenas OR ((oat OR oats) NEAR/2 (plant* OR hull* OR bran* OR kernel* OR groat*)))) AND DT=(article)
Cochrane (metabol* OR cometabol* OR co-metabol* OR nutrient* OR nutraceut* OR phytochem* OR phytopharmaceut* OR pharmacokinetic* OR ((dietary OR nutritional) NEAR/3 (supplement*)) OR "Amino acids" OR "Amino acid" OR "Cyanogenic glucoside" OR "Cyanogenic glucosides" OR "Fatty acids" OR "Fatty acid" OR "Nucleic acid base" OR "Nucleic acid bases" OR "Nucleic acid based" OR "Organic acid" OR "Organic acids" OR "Organosulfur compound" OR "Organosulfur compounds" OR "Phenolic acids" OR "Phenolic acid" OR "Sugar alcohol" OR "Sugar alcohols" OR Alkaloid* OR Alkane* OR Amine* OR Benzenoid* OR Carotenoid* OR Chlorophyll* OR Disaccharide* OR Flavonoid* OR Indole* OR Lignan* OR Monosaccharide* OR Peptide* OR Polyacetylene* OR Polysaccharide* OR Protein* OR Purine* OR Pyrimidine* OR Stilbene* OR Terpene* OR Terpenoid* OR avenanthramid* OR amide*) AND (chromatogra* OR electrochromatogra* OR ((plant OR plants) NEAR/3 (extract* OR biochem* OR preparation* OR medicinal*))) AND (buckwheat* OR fagopyrum "avena sativa" OR "avena nuda" OR avena OR avenas OR ((oat OR oats) NEAR/3 (plant* OR hull* OR bran* OR kernel* OR groat*)))
PubMed ((((("Metabolism"[mh] OR "Metabolomics"[mh] OR "Energy Metabolism"[mh] OR "Phytochemicals"[mh] OR "Nutrients"[mh] OR "Dietary Supplements"[mh] OR "Pharmacokinetics"[mh] OR metabol*[tiab] OR cometabol*[tiab] OR co-metabol*[tiab] OR nutrient*[tiab] OR nutraceut*[tiab] OR phytochem*[tiab] OR phytopharmaceut*[tiab] OR pharmacokinetic*[tiab] OR ((dietary[tiab] OR nutritional[tiab]) AND (supplement*[tiab]))) OR ("Alkaloids"[mh] OR "Amines"[mh] OR "Amino acids"[mh] OR "Carbohydrates"[mh] OR "Carotenoids"[mh] OR "Disaccharides"[mh] OR "Fatty Acids"[mh] OR "Flavonoids"[mh] OR "Indoles"[mh] OR "Lignans"[mh] OR "Lipids"[mh] OR "Monosaccharides"[mh] OR "Peptides"[mh] OR "Polysaccharides"[mh] OR "Proteins"[mh] OR "Purines"[mh] OR "Pyrimidines"[mh] OR "Stilbenes"[mh] OR "Sugar Alcohols"[mh] OR "Terpenes"[mh] OR "Amides"[mh] OR Amino acid*[tiab] OR Cyanogenic glucoside*[tiab] OR Fatty acid*[tiab] OR Nucleic acid base*[tiab] OR Organic acid*[tiab] OR Organosulfur compound*[tiab] OR Phenolic acid*[tiab] OR Sugar alcohol*[tiab] OR Alkaloid*[tiab] OR Alkane*[tiab] OR Amine*[tiab] OR Benzenoid*[tiab] OR Carotenoid*[tiab] OR Chlorophyll*[tiab] OR Disaccharide*[tiab] OR Flavonoid*[tiab] OR Indole*[tiab] OR Lignan*[tiab] OR Monosaccharide*[tiab] OR Peptide*[tiab] OR Polyacetylene*[tiab] OR Polysaccharide*[tiab] OR Protein[tiab] OR Proteins[tiab] OR Purine*[tiab] OR Pyrimidine*[tiab] OR Stilbene*[tiab] OR Terpene*[tiab] OR Terpenoid*[tiab] OR avenanthramide*[tiab] OR amide*[tiab])) AND (("Plant Extracts"[mh] OR "Chromatography, Liquid"[Mesh] OR chromatogra*[tiab] OR electrochromatogra*[tiab] OR ((plant[tiab] OR plants[tiab]) AND (extract*[tiab] OR biochem*[tiab] OR preparation*[tiab] OR medicinal*[tiab]))) AND ("Fagopyrum"[mh] OR buckwheat*[tiab] OR fagopyrum[tiab] OR "Avena"[mh] OR avena sativa[tiab] OR avena nuda[tiab] OR avena[tiab] OR avenas[tiab] OR ((oat[tiab] OR oats[tiab]) AND (plant[tiab] OR plants[tiab] OR hull*[tiab] OR bran[tiab] OR brans[tiab] OR kernel*[tiab] OR groat*[tiab]))) NOT (letter[pt] OR news[pt] OR comment[pt] OR editorial[pt] OR congress[pt]))

Supplemental table 2. Flavonoids, phenolic compounds, AVAs in oat and buckwheat				
Flavonoids in oat and buckwheat				
Compound	Plant and specie (or cultivar)	Plant part	Concentration	Publication
Total flavonoids	Tartary Buckwheat	Seeds	0.037 %	(Jiang, <i>et al.</i> , 2007)
	Buckwheat	Whole grain	7mg RE/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	Hull	17mg RE/g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	groat	6 mg RE/100g	(Sedej, <i>et al.</i> , 2012)
	Common Buckwheat	Seeds	2.038 %	(Jiang, <i>et al.</i> , 2007)
	Common Buckwheat	Sprouts	507.6 mg RE/ 100g	(Chen, <i>et al.</i> , 2020)
	Common Buckwheat	Seed and hull	371.5- 1463.7 mg/100g	(Oomah & Mazza, 1996)
	Common buckwheat	groats	170 mg RE/100g	(Tien, <i>et al.</i> , 2018)
	Tartary buckwheat	groats	1434 mg RE/100g	(Tien, <i>et al.</i> , 2018)
Isoorientin	Common Buckwheat	Grain	0.15-7.61 mg/100g	(Terpinc, <i>et al.</i> , 2016)
	Common Buckwheat	Sprouts, microgreens, leafy greens	751.53 mg/100g	(Sharma, 2011)
Orientin	Common Buckwheat	Sprouts, microgreens, leafy greens	353.25 mg/100g	(Sharma, 2011)
Apigenin	Common Buckwheat	Grain	0.05-0.24 mg/100g	(Terpinc, <i>et al.</i> , 2016)
Fagopyrin	Common and Tartary Buckwheat	Leaves, flowers	19-32 mg/100g	(Habtemariam, 2019)
	Tartary buckwheat	All parts	3.06- 144.6 mg/100g	(Kočevár Glavač, <i>et al.</i> , 2017)
	Common Buckwheat	Sprouts	0.0025 to 0.041 % of dry mass	(Kreft, 2013)
	Tartary Buckwheat		0.10 to 0.12 %.	(Kreft, 2013)
Kaempferol-3-O-rutinoside	Buckwheat Daesan maemil	Seeds	3.3 mg/100g	(Lee, <i>et al.</i> , 2013)
	Buckwheat Yangjul maemil	Seeds	3.2 mg/100g	(Lee, <i>et al.</i> , 2013)
	Tartary Buckwheat	Seeds	8.2 mg/100g	(Lee, <i>et al.</i> , 2013)
Kaempferol	Tartary Buckwheat Chuanqiao 1 &2, Xiqiao 3	flour	1-3 mg/100g	(Peng L., 2017)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	bran	2-4 mg/100g	(Peng L., 2017)
	Buckwheat	Hull, Groat	0.02 mg/100g	(Dziedzic, <i>et al.</i> , 2009)
	Tartary Buckwheat	Flowers, Leaves, Stalk, roots	0.14-0.33 mg/100g	(Dziedzic, <i>et al.</i> , 2018)
Quercetin	Oat (Mina DS)	germ, endosperm	2.4-8.9 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat (Mina DS)	husks	3.76-8.82 mg/100g	(Kerlene, <i>et al.</i> , 2015)
	Oat (Migla DS)	germ, endosperm	1.22-5.16 mg/100g	(Kerlene, <i>et al.</i> , 2015)
	Oat (Migla DS)	husks	0.27-4.83 mg/100g	(Kerlene, <i>et al.</i> , 2015)
	Buckwheat	grain with husks	0.31-0.67 mg/100g	(Kerlene, <i>et al.</i> , 2015)
	Buckwheat	husks	0.28–0.41 mg/100g	(Kerlene, <i>et al.</i> , 2015)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	flour	24-72 mg/100g	(Peng L., 2017)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	bran	62-111 mg/100g	(Peng L., 2017)
	Tartary Buckwheat (Kitawase, Hokkai T8, T9, T10)	Sprouts	1.3-47 mg/100g	(Arasu, <i>et al.</i> , 2014)
	Common Buckwheat/ Tartary Buckwheat	Bran	12.7-13.0 mg/100g	(Bai, <i>et al.</i> , 2015)
	Buckwheat	Hull, Groat	0.02-0.59 mg/100g	(Dziedzic, <i>et al.</i> , 2009)
	Tartary Buckwheat	Flowers, leaves, Stalk, roots	0.21-84.5 mg/100g	(Dziedzic, <i>et al.</i> , 2018)
	Common Buckwheat	Hull	0.02-0.07 mg/100g	(Kalinova, <i>et al.</i> , 2019)
	Common Buckwheat/ Tartary Buckwheat (Kitawase, Hokkai T9)	Seeds, Sprouts	10 mg/100g	(Kim, <i>et al.</i> , 2008)
	Common Buckwheat	Hull	0.61 mg/100g	(Quettier-Deleu, <i>et al.</i> , 2000)
	Common Buckwheat	Flour	0.15 mg/100g	(Quettier-Deleu, <i>et al.</i> , 2000)
	Common Buckwheat	Sprouts	33 mg/100 g	(Chen, <i>et al.</i> , 2020)
	Tartary buckwheat	Seed	425.65-857.62mg/100g	{Guo, 2011 #156}

	Tartary Buckwheat	Bran	330 mg/100g	(Ge & Wang, 2020)
Quercetin	Common buckwheat	Hulls, flour	0.15-0.61 mg/100g	(Quettier-Deleu, <i>et al.</i> , 2000)
	Tartary Buckwheat	All parts	0-82 mg/100g	(Kočevar Glavač, <i>et al.</i> , 2017)
	Common Buckwheat	Grain	1.19-7.69 mg/100g	(Terpinc, <i>et al.</i> , 2016)
	Common Buckwheat	Sprouts, microgreens, leafy greens	17.12 mg/100g	(Sharma, 2011)
	Tartary buckwheat	Groats	116 mg/100g	(Tien, <i>et al.</i> , 2018)
Hyperoside	Common buckwheat	Hulls, flour	0.2-1.6 mg/100g	(Quettier-Deleu, <i>et al.</i> , 2000)
Luteolin	Common Buckwheat	Grain	0.33-4.13 mg/100g	(Terpinc, <i>et al.</i> , 2016)
Kaempferol	Common Buckwheat	Grain	0.1-0.29 mg/100g	(Terpinc, <i>et al.</i> , 2016)
Naringenin	Common Buckwheat	Grain	0.13-0.60 mg/100g	(Terpinc, <i>et al.</i> , 2016)
Quercitrin	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	flour	43-68 mg/100g	(Peng L., 2017)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	bran	146-291 mg/100g	(Peng L., 2017)
Rutin	Buckwheat	Grain with husks	14.7–40.6 mg/100g	(Kerlene, <i>et al.</i> , 2015)
	Common Buckwheat	Seed and hull	44.2-85.3 mg/100g	(Oomah & Mazza, 1996)
	Buckwheat	husks	11.3–13.98 mg/100g	(Kerlene, <i>et al.</i> , 2015)
	Husked oat	Grain without husks	n.d.	(Kerlene, <i>et al.</i> , 2015)
	Husked oat	husks	0.00–3.02 mg/100g	(Kerlene, <i>et al.</i> , 2015)
	Oat	naked	0.22-0.47 mg/100g	{Tong, 2014 #155}
	Common oat	Grain without husks	n.d.	(Kerlene, <i>et al.</i> , 2015)
	Common oat	husks	n.d.	(Kerlene, <i>et al.</i> , 2015)
	Tartary Buckwheat Ishisoba	grain	1193 mg/100g	(Andrea Brunori, 2009)
	Tartary Buckwheat golden	grain	1041mg/100g	(Andrea Brunori, 2009)
	Tartary Buckwheat Donan	grain	979 mg/100g	(Andrea Brunori, 2009)
	Common buckwheat	hull	80-440 mg/100g	(Steadman, <i>et al.</i> , 2001)
	Common Buckwheat	Sprouts, microgreens, leafy greens	1440.92 mg/100g	(Sharma, 2011)
	Common Buckwheat	Bran	20-30 mg/100g	(Steadman, <i>et al.</i> , 2001)
	Common Buckwheat	groat	18 mg/100g	(Steadman, <i>et al.</i> , 2001)
	Tartary Buckwheat	groat	84 mg/100g	(Steadman, <i>et al.</i> , 2001)
	Common Buckwheat	hulls	84 mg/100g	(Steadman, <i>et al.</i> , 2001)
	Tartary Buckwheat	hulls	437 mg/100g	(Steadman, <i>et al.</i> , 2001)
	Buckwheat	whole grain	14.6 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	hull	22.5 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	groat	11.6 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat Daesan maemil	Seeds	45.0 mg/100g	(Lee, <i>et al.</i> , 2013)
	Buckwheat Yangjul maemil	Seeds	26.6 mg/100g	(Lee, <i>et al.</i> , 2013)
	Tartary Buckwheat	Seeds	204.0 mg/100g	(Lee, <i>et al.</i> , 2013)
	Tartary Buckwheat	seeds	810–1700 mg/100g	(Fabjan, <i>et al.</i> , 2003)
	Common Buckwheat	seeds	10mg/100g	(Fabjan, <i>et al.</i> , 2003)
	Tartary Buckwheat	Flower, leaf, Stalk	52.38-294.9 mg/100g	(Dziedzic, <i>et al.</i> , 2018)
	Common/Tartary Buckwheat	Flower. leaf, stalk, seed	0.42-7.77% <sup>a</sup>	(Zielińska, <i>et al.</i> , 2012)
	Tartary Buckwheat (Kitawase, Hokkai T8, T9, T10)	Sprouts	329-4793 mg/100g	(Arasu, <i>et al.</i> , 2014)
	Tartary Buckwheat/ Common Buckwheat	Bran	0.33 mg/100g	(Bai, <i>et al.</i> , 2015)
	Common Buckwheat/Tartary Buckwheat (Kitawase, Hokkai T9)	Seeds, sprouts	20-2380 mg/100g	(Kim, <i>et al.</i> , 2008)
	Common Buckwheat	Sprouts	719 mg/ 100g	(Chen, <i>et al.</i> , 2020)
	Common Buckwheat	Flour	12.7 mg/100g	(Danila, <i>et al.</i> , 2007)
	Buckwheat	Hull, Groat	0.7-3.59 mg/100g	(Dziedzic, <i>et al.</i> , 2009)
	Common Buckwheat	Seeds	1400-18850 mg/100g	(A. R. Gulpinar, <i>et al.</i> , 2012)
	Common Buckwheat	Hull	0.76-0.98 mg/100g	(Kalinova, <i>et al.</i> , 2019)
	Common Buckwheat	Hull	5.21 mg/100g	(Quettier-Deleu, <i>et al.</i> , 2000)
	Common Buckwheat	Flour	2.28 mg/ 100g	(Quettier-Deleu, <i>et al.</i> , 2000)
	Husked oat	husks	0.00-3.02 mg/100g	(Kerlene, <i>et al.</i> , 2015)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Flour	849-1486 mg/100g	(Peng L., 2017)
	Tartary buckwheat	Seed	518.54-1325.59 mg/100g	{Guo, 2011 #156}
	Tartary Buckwheat	Bran	4079-5186mg/100g	(Peng L., 2017)

	Chuanqiao 1&2, Xiqiao 3			
	Tartary Buckwheat	Bran	332.9 mg/100g	(Ge & Wang, 2020)
	Common Buckwheat	Leaves, flowers	822-3417 mg/100g	(Habtemariam, 2019)
	Common Buckwheat	Flowers	568 mg/100g	(Kraujaliene, <i>et al.</i> , 2017)
	Common Buckwheat	Seeds	54.2 mg/ 100g	(Park, <i>et al.</i> , 2019)
Rutin	Tartary Buckwheat	All parts	157-3245mg/100g	(Kočevlar Glavač, <i>et al.</i> , 2017)
	Common Buckwheat	Grains	0.25-33.0 mg/100 g	(Hung & Morita, 2008)
	Common buckwheat	Hulls, flour	2.28- 5.205 mg/100g	(Quettier-Deleu, <i>et al.</i> , 2000)
Epicatechin	Common buckwheat	Hulls, flour	1.15-3.4 mg/100g	(Quettier-Deleu, <i>et al.</i> , 2000)
Epicatechin gallate	Common buckwheat	Hulls, flour	0.88 mg/100g	(Quettier-Deleu, <i>et al.</i> , 2000)
Catechin	Tartary buckwheat	Seed	8.89-19.96 mg/100g	{Guo, 2011 #156}
Phenolics				
Compound	Plant	Plant part	Concentration	Author
Total phenolics	Oat (Mina DS)	Germ, endosperm	180 mg RE/100g <sup>a</sup>	(Keriene, <i>et al.</i> , 2015)
	Oat (Mina DS)	Hull (or husks)	265 mg RE/100g <sup>a</sup>	(Keriene, <i>et al.</i> , 2015)
	Oat (Migla DS)	Germ, endosperm	286 mg RE/100g <sup>a</sup>	(Keriene, <i>et al.</i> , 2015)
	Oat (Migla DS)	Hull (or husks)	576 mg RE/100g <sup>a</sup>	(Keriene, <i>et al.</i> , 2015)
	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	35.1-87.4 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat (Ogle)	Germ, endosperm, bran, hull	23.2 mg/100g	(Xing & White, 1997)
	Oat (Ogle)	Hull	35.1 mg/100g	(Xing & White, 1997)
	Oat (Wild oat, SH430, M73)	Germ, endosperm, bran, hull	0.14-0.16 mg (per seed)	(Gallagher, <i>et al.</i> , 2010)
	Oat (13 cultivars)	Germ, endosperm, bran, hull	3.57-14.35 mg/100g	(Chen, <i>et al.</i> , 2018)
	Oat (8 cultivars)	Germ, endosperm, bran, hull	120.1-168.4 mg/100g	(Multari, <i>et al.</i> , 2018)
	Common Buckwheat	grain with husks	1020 mg RE/100g	(Keriene, <i>et al.</i> , 2015)
	Common Buckwheat	husks	1410 mg RE/100g	(Keriene, <i>et al.</i> , 2015)
	Oat (commercial)	Bran	2.52 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (commercial), thermally processed	Bran	3.17 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (22 commercial varieties)	Seeds and bran	64.5-151.8 mg/100 g	(Soycan, <i>et al.</i> , 2019)
	Oat		113.8 mg GAE/100g	(Holasovaa, <i>et al.</i> , 2002)
	Buckwheat	seeds	330.3 GAE/100g	(Holasovaa, <i>et al.</i> , 2002)
	uckwheat	Dehulled seeds	390.3 GAE/100g	(Holasovaa, <i>et al.</i> , 2002)
	uckwheat	leaves	3.951.4 mg GAE/100g	(Holasovaa, <i>et al.</i> , 2002)
	Common Buckwheat	groat	244 mg/100g****	(Steadman, <i>et al.</i> , 2001)
	Common Buckwheat	whole grain	142 mg/100g***	(Sedej, <i>et al.</i> , 2012)
	Common Buckwheat	hull	220 mg/100g***	(Sedej, <i>et al.</i> , 2012)
	Common Buckwheat	groat	141 mg/100g***	(Sedej, <i>et al.</i> , 2012)
	Common Buckwheat	Seed milling fractions	1548 mg/100g****	(Steadman, <i>et al.</i> , 2001)
	Common Buckwheat	Seed, steam, aerial parts	9030-30100 mg/100g***	(A. R. Gulpinar, <i>et al.</i> , 2012)
	Common Buckwheat	Sprouts	778.6 mg/ 100g***	(Chen, <i>et al.</i> , 2020)
	Tartary Buckwheat	Bran, flour	123 mg RE/ 100g	(Ge & Wang, 2020)
	Common Buckwheat	Seeds	537 mg GAE/ 100g	(Liu, <i>et al.</i> , 2019)
	Tartary Buckwheat	Seeds	667 mg GAE/100g	(Liu, <i>et al.</i> , 2019)
	Common Buckwheat	Seeds	177.7 mg/100g	(Rocchetti, <i>et al.</i> , 2019)
	Common Buckwheat	Seeds	1495 mg/100g	(Siwatch, <i>et al.</i> , 2019)
	Common buckwheat	Seeds, Sprouts	64.5-670 mg/100g**	(Alvarez-Jubete, <i>et al.</i> , 2010)
	Common buckwheat	groats	773 mg FAE/100g	(Tien, <i>et al.</i> , 2018)
	Tartary buckwheat	groats	1927 mg FAE/100g	(Tien, <i>et al.</i> , 2018)
	Common buckwheat	Hull, seed, dehulled seed	280- 513.18 chlorogenic acid equivalents/100g	(Dziadek, <i>et al.</i> , 2016)
	Oat	naked	101.7 to 151.9 mg/100g	{Tong, 2014 #155}
	Common buckwheat	hulls, brans and flour	1386-2487 GAE mg/100g	(Li, <i>et al.</i> , 2013)
4-hydrobenzoic acid	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	0.62-5.06 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat (Wild oat, SH430, M73)	Germ, endosperm, bran, hull	0.0004-0.0007 mg (per seed)	(Gallagher, <i>et al.</i> , 2010)
	Oat (13 cultivars)	Germ, endosperm, bran, hull	0.67-2.37 mg/100g	(Chen, <i>et al.</i> , 2018)
	Oat (22 commercial varieties)	Seeds and bran	0.2-1.55 mg/100 g	(Soycan, <i>et al.</i> , 2019)
	Oat	bran	2.2 mg/100g	(Mattila, <i>et al.</i> , 2005)
	Oat (20 varieties)	Seeds	0.54 mg/100g	(Varga, <i>et al.</i> , 2018)
	Oat (20 varieties)	Hull	2.25 mg/100g	(Varga, <i>et al.</i> , 2018)
	Tartary Buckwheat	Flowers, leaves, stalk,	0.07-0.14 mg/100g	(Dziedzic, <i>et al.</i> , 2018)

		roots		
	Tartary buckwheat	Seed	2.22- 8.78 mg/100g	{Guo, 2011 #156}
gallic acid	Buckwheat	Hull, Groat	0.02-0.17 mg/100g	(Dziedzic, <i>et al.</i> , 2009)
	Tartary Buckwheat	Flower, leaf, root	0.08-0.73 mg/100g	(Dziedzic, <i>et al.</i> , 2018)
	Oat (13 cultivars)	Germ, endosperm, bran, hull	1.43-7.05 mg/100g	(Chen, <i>et al.</i> , 2018)
	Common Buckwheat	Seeds	0.28 mg/100g	(Liu, <i>et al.</i> , 2019)
	Tartary Buckwheat	Seeds	0.65 mg/100g	(Liu, <i>et al.</i> , 2019)
	Common Buckwheat	Seeds	0.98 mg/100g	(Park, <i>et al.</i> , 2019)
	Buckwheat	seed	0.48-0.62 mg/100g	{Guo, 2011 #156}
	Common Buckwheat	Grain	0.07-0.19 mg/100g	(Terpinc, <i>et al.</i> , 2016)
Chlorogenic acid	Tartary Buckwheat	Flower, leaf, root	0.21-101.34 g/100g	(Dziedzic, <i>et al.</i> , 2018)
	Oat	naked	0.20 mg/100g	{Tong, 2014 #155}
	Tartary Buckwheat (Kitawase, Hokkai T8, T9, T10)	Sprouts	162-377mg/100g	(Arasu, <i>et al.</i> , 2014)
	Tartary Buckwheat	Flowers, Leaves, stalk, roots	2.13-101 mg/100g	(Dziedzic, <i>et al.</i> , 2018)
	Common Buckwheat/Tartary Buckwheat (Kitawase, Hokkai T9)	Seeds, sprouts	80-170 mg/100g	(Kim, <i>et al.</i> , 2008)
	Common Buckwheat	Seeds	0.26-1.26 mg/100g	(Kiprovski, <i>et al.</i> , 2015)
	Common Buckwheat	Sprouts, microgreens, leafy greens	156.23 mg/100g	(Sharma, 2011)
	Common Buckwheat	Flowers	124.7 mg/100g	(Kraujaliene, <i>et al.</i> , 2017)
	Common Buckwheat	Seeds	1.15 mg/100g	(Liu, <i>et al.</i> , 2019)
	Tartary Buckwheat	Seeds	0.57 mg/ 100g	(Liu, <i>et al.</i> , 2019)
	Common Buckwheat	Seeds	11.0 mg/100g	(Park, <i>et al.</i> , 2019)
	Common and tartary buckwheat		3.1-16.6 mg/100g	(Syta, 2014)
vanillic acid	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	0.39-9.38 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat (Mina DS)	Germ, endosperm	0.26 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat	bran	0.4 mg/100g	(Mattila, <i>et al.</i> , 2005)
	Oat (Mina DS)	Hull (or husks)	1.12 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat (Migla DS)	Germ, endosperm	0.34 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat (Migla DS)	Hull (or husks)	0.89 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat (Ogle)	Germ, endosperm, bran, hull	0.34 mg/100g	(Xing & White, 1997)
	Oat (Ogle)	Hull	5.42 mg/100g	(Xing & White, 1997)
	Oat	naked	0.51 mg/100g	{Tong, 2014 #155}
	Oat (Wild oat, SH430, M73)	Germ, endosperm, bran, hull	0.0007-0.0014 mg (per seed)	(Gallagher, <i>et al.</i> , 2010)
	Oat (13 cultivars)	Germ, endosperm, bran, hull	037-4.75 mg/100g	(Chen, <i>et al.</i> , 2018)
	Oat (unspecified)	hull	593 mg/100g (fermented)	(Garleb, <i>et al.</i> , 1991)
	Oat (8 cultivars)	Germ, endosperm, bran, hull	0.46-0.71 mg/100g	(Multari, <i>et al.</i> , 2018)
	Oat (commercial)	Bran	1.01 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (commercial), thermally processed	Bran	1.25 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (22 commercial varieties)	Seeds and bran	0.49-1.29 mg/100 g	(Soycan, <i>et al.</i> , 2019)
	Oat (20 varieties)	Seeds	0.81 mg/100g	(Varga, <i>et al.</i> , 2018)
	Oat (20 varieties)	Hull	20.6 mg/100g	(Varga, <i>et al.</i> , 2018)
	Buckwheat	whole grain	1.5 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	hull	3.71 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Common Buckwheat	Hull	0.03-0.05 mg/100g	(Kalinova, <i>et al.</i> , 2019)
	Buckwheat	seed	0.21-1.6 mg/100g	{Guo, 2011 #156}
	Common and tartary buckwheat		221.1- 311.7 mg/100g	(Syta, 2014)
syringic acid	Buckwheat	whole grain	5.23 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	hull	3.63 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	groat	6.35 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Oat	bran	2.8 mg/100g	(Mattila, <i>et al.</i> , 2005)
	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	0.46-0.79 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat (Mina DS)	Germ, endosperm	0.34 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat (Mina DS)	Hull (or husks)	0.68 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat (Migla DS)	Germ, endosperm	0.55 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)

	Oat (Migla DS)	Hull (or husks)	0.43 mg/100g <sup>a</sup>	(Kerienne, <i>et al.</i> , 2015)
	Oat (8 cultivars)	Germ, endosperm, bran, hull	0.39-0.89 mg/100g	(Multari, <i>et al.</i> , 2018)
	Oat (22 commercial varieties)	Seeds and bran	0.45-2.11 mg/100 g	(Soycan, <i>et al.</i> , 2019)
	Oat (20 varieties)	Seeds	0.49 mg/100g	(Varga, <i>et al.</i> , 2018)
	Oat (20 varieties)	Hull	3.88 mg/100g	(Varga, <i>et al.</i> , 2018)
	Tartary Buckwheat	Flower, leaf, root	0.12-0.56mg/100g	(Dziedzic, <i>et al.</i> , 2018)
	Buckwheat	seed	0.12-0.18mg/100g	{Guo, 2011 #156}
	Common Buckwheat	Seeds	0.37 mg/ 100g	(Liu, <i>et al.</i> , 2019)
syngaldehyde	Tartary Buckwheat	Seeds	0.19 mg/100g	(Liu, <i>et al.</i> , 2019)
	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	0.32-2.12 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat (unspecified)	hull	566 mg/100g (fermented)	(Garleb, <i>et al.</i> , 1991)
caffeic acid	Oat (8 cultivars)	Germ, endosperm, bran, hull	0.11-0.17 mg/100g	(Multari, <i>et al.</i> , 2018)
	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm	0.52-1.57 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat (Ogle)	Germ, endosperm, bran, hull	1.68 mg/100g	(Xing & White, 1997)
	Oat (Ogle)	Hull	-	(Xing & White, 1997)
	Oat (13 cultivars)	Germ, endosperm, bran, hull	0.09-0.70 mg/100g	(Chen, <i>et al.</i> , 2018)
	Oat (commercial)	Bran	0.86 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (commercial), thermally processed	Bran	2.28 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat	grains	0.2-23.9 nmol/g	(Skoglund, <i>et al.</i> , 2008)
	Oat (22 commercial varieties)	Seeds and bran	1.04-7.27 mg/100 g	(Soycan, <i>et al.</i> , 2019)
	Oat (20 varieties)	Seeds	0.86 mg/100g	(Varga, <i>et al.</i> , 2018)
	Oat (20 varieties)	Hull	0.42 mg/100g	(Varga, <i>et al.</i> , 2018)
	Common Buckwheat	fiber	1.12 mg/100g	(Wefers & Bunzel, 2015)
	Tartary Buckwheat	Flower, leaf, root	0.29-1.84 mg/100g	(Dziedzic, <i>et al.</i> , 2018)
	Buckwheat	seed	0.12-0.49 mg/100g	{Guo, 2011 #156}
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	flour	22-47 mg/100g	(Peng L., 2017)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	bran	65-119 mg/100g	(Peng L., 2017)
	Common Buckwheat	Seeds	9.11 mg/ 100g	(Park, <i>et al.</i> , 2019)
	Oat	bran	0.5 mg/100g	(Mattila, <i>et al.</i> , 2005)
	Common buckwheat	Seeds, sprouts	8.8-15.1 umol/100g	(Alvarez-Jubete, <i>et al.</i> , 2010)
	Buckwheat	Hull, Groat	0.04 mg/100g	(Dziedzic, <i>et al.</i> , 2009)
	Tartary Buckwheat	Flowers, leaves, stalk, roots	0.29-1.84 mg/100g	(Dziedzic, <i>et al.</i> , 2018)
	Common Buckwheat	Grains	0.12-3.34 mg/100g	(Terpinc, <i>et al.</i> , 2016)
sinapic acid	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	0.91-1.03 mg/100g <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat	bran	9 mg/100g	(Mattila, <i>et al.</i> , 2005)
	Oat (Mina DS)	Germ, endosperm	2.4 mg/100g <sup>a</sup>	(Kerienne, <i>et al.</i> , 2015)
	Oat (Mina DS)	Hull (or husks)	0.14 mg/100g <sup>a</sup>	(Kerienne, <i>et al.</i> , 2015)
	Oat (Migla DS)	Germ, endosperm	2.81 mg/100g <sup>a</sup>	(Kerienne, <i>et al.</i> , 2015)
	Oat (Migla DS)	Hull (or husks)	0.27 mg/100g <sup>a</sup>	(Kerienne, <i>et al.</i> , 2015)
	Oat (Ogle)	Hull	0.56 mg/100g	(Xing & White, 1997)
	Oat (commercial)	Bran	0.47 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (commercial), thermally processed	Bran	0.53 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (22 commercial varieties)	Seeds and bran	1.89-8.03 mg/100 g	(Soycan, <i>et al.</i> , 2019)
	Buckwheat	Hull, Groat	0.41 mg/100g	(Dziedzic, <i>et al.</i> , 2009)
	Buckwheat	whole grain	2.81 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	hull	2.98 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	groat	3.04 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Common Buckwheat	fibre	3.24 mg/100g	(Wefers & Bunzel, 2015)
	Common Buckwheat	Seeds	0.35 mg/100g	(Liu, <i>et al.</i> , 2019)
ferulic acid	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	0.66-0.84 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat (Mina DS)	Germ, endosperm	9.87 mg/100g <sup>a</sup>	(Kerienne, <i>et al.</i> , 2015)
	Oat	Grain	16.50-149.36 mg/100g	(Kovacova & MaliNová, 2007)

	Oat	naked	0.23 mg/100g	{Tong, 2014 #155}
	Oat (Mina DS)	Hull (or husks)	30.9 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat	bran	14 mg/100g	(Mattila, <i>et al.</i> , 2005)
	Oat	grain	0.07-0.6 mg/100g	(Dokuyucu, <i>et al.</i> , 2003)
	Oat (Migla DS)	Germ, endosperm	10.5 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat (Migla DS)	Hull (or husks)	45.85 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat (Ogle)	Germ, endosperm, bran, hull	14.7 mg/100g	(Xing & White, 1997)
	Oat (Ogle)	Hull	14.2 mg/100g	(Xing & White, 1997)
	Oat (Wild oat, SH430, M73)	Germ, endosperm, bran, hull	0.09-0.13 mg (per seed)	(Gallagher, <i>et al.</i> , 2010)
	Oat (13 cultivars)	Germ, endosperm, bran, hull	0.14-1.89 mg/100g	(Chen, <i>et al.</i> , 2018)
	Oat (unspecified)	hull	214 mg/100g (fermented)	(Garleb, <i>et al.</i> , 1991)
	Oat (8 cultivars)	Germ, endosperm, bran, hull	53.2-82.9 mg/100g	(Multari, <i>et al.</i> , 2018)
	Oat (commercial)	Bran	0.46 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (commercial), thermally processed	Bran	0.49 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (22 commercial varieties)	Seeds and bran	15.2-115.3 mg/100g	(Soycan, <i>et al.</i> , 2019)
	Oat (20 varieties)	Seeds	32.7 mg/100g	(Varga, <i>et al.</i> , 2018)
	Oat (20 varieties)	Hull	809.5 mg/100g	(Varga, <i>et al.</i> , 2018)
	Oat	bran	33.0 mg/100g	(Mattila, <i>et al.</i> , 2005)
	Buckwheat	whole grain	1.72 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	hull	1.75 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	groat	1.74 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Common and tartary buckwheat		5.6-7.3 mg/100g	(Syta, 2014)
	Common Buckwheat	fibre	4.42 mg/100g	(Wefers & Bunzel, 2015)
	Tartary Buckwheat	Flower, leaf, root	0.32-1.18 mg/100g	(Dziedzic, <i>et al.</i> , 2018)
	Common Buckwheat	Seeds	0.89 mg/100g	(Liu, <i>et al.</i> , 2019)
	Common Buckwheat	Grains	0.17-42.8 µg/g	(Hung & Morita, 2008)
	Buckwheat	seed	1.86-7.29mg/100g	{Guo, 2011 #156}
Cis-ferulic acid	Common Buckwheat	fibre	0.93 mg/100g	(Wefers & Bunzel, 2015)
Trans-ferulic acid	Common and tartary buckwheat		31.7-65.7mg/100g	(Syta, 2014)
p-anisic	Common and tartary buckwheat		744.5-1190 mg/100g	(Syta, 2014)
Methoxy-cinnamic acid	Common and tartary buckwheat		14.1-74.7 mg/100g	(Syta, 2014)
p -coumaric acid	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	0.24-1.19 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat	bran	1.2 mg/100g	(Mattila, <i>et al.</i> , 2005);
	Oat	grain	0.12-0.19 mg/100g	(Dokuyucu, <i>et al.</i> , 2003)
	Oat (Mina DS)	Germ, endosperm	0.57 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat (Mina DS)	Hull (or husks)	26.9 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat (Migla DS)	Germ, endosperm	1.13 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat (Migla DS)	Hull (or husks)	39.9 mg/100g <sup>a</sup>	(Kerlene, <i>et al.</i> , 2015)
	Oat	grains	0.3-29.7 nmol/g	(Skoglund, <i>et al.</i> , 2008)
	Oat (Ogle)	Germ, endosperm, bran, hull	4.49 mg/100g	(Xing & White, 1997)
	Oat (Ogle)	Hull	59.7 mg/100g	(Xing & White, 1997)
	Oat (Wild oat, SH430, M73)	Germ, endosperm, bran, hull	0.02-0.04 mg (per seed)	(Gallagher, <i>et al.</i> , 2010)
	Oat (unspecified)	hull	245 mg/100g (fermented)	(Garleb, <i>et al.</i> , 1991)
	Oat (8 cultivars)	Germ, endosperm, bran, hull	59.0-82.6 mg/100g	(Multari, <i>et al.</i> , 2018)
	Oat (commercial)	Bran	0.26 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (commercial), thermally processed	Bran	0.30 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (22 commercial varieties)	Seeds and bran	0.19-4.94 mg/100 g	(Soycan, <i>et al.</i> , 2019)
	Oat (20 varieties)	Seeds	6.22 mg/100g	(Varga, <i>et al.</i> , 2018)
	Oat (20 varieties)	Hull	368 mg/100g	(Varga, <i>et al.</i> , 2018)
	Common Buckwheat	fibre	3.01 mg/100g	(Wefers & Bunzel, 2015)
	Buckwheat	seed	0.23-0.68mg/100g	{Guo, 2011 #156}
	Tartary Buckwheat	Flower, leaf, root	0.08-1.69 mg/100g	(Dziedzic, <i>et al.</i> , 2018)
	Buckwheat	Hull, Groat	0.07-0.29 mg/100g	(Dziedzic, <i>et al.</i> , 2009)
Cis- p-coumaric acid	Common Buckwheat	fibre	0.589 mg/100g	(Wefers & Bunzel, 2015)
coumaric	Oat	grain	8.05-210.27 mg/100 g	(Kovacova & MaliNová,



				2007)
Ferulic acid	Oat	grains	0.4-15.4 nmol/l	(Skoglund, <i>et al.</i> , 2008)
o-coumaric acid	Oat (Ogle)	Hull	0.69 mg/100g	(Xing & White, 1997)
	Oat (8 cultivars)	Germ, endosperm, bran, hull	0.11-0.38 mg/100g	(Multari, <i>et al.</i> , 2018)
2-hydroxycinnamic acid	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	0.05-0.67 mg/100g	(Shewry, <i>et al.</i> , 2008)
2,4-dihydrobenzoic acid (Protocatechuic acid )	Oat (13 cultivars)	Germ, endosperm, bran, hull	0.10-1.04 mg/100g	(Chen, <i>et al.</i> , 2018)
	Buckwheat	whole grain	9.26 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	hull	16.8 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	groat	10.3 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat	seed	1.71-4.63mg/100g	{Guo, 2011 #156}
	Common Buckwheat	Hull	0.18-0.21 mg/100g	(Kalinova, <i>et al.</i> , 2019)
	Common Buckwheat	Seeds	0.4 mg/100g	(Liu, <i>et al.</i> , 2019)
	Tartary Buckwheat	Seeds	1.0 mg/100g	(Liu, <i>et al.</i> , 2019)
	Oat (Cacko)	Germ, endosperm, bran, hull	0.50 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat (commercial)	Bran	5.50 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (commercial), thermally processed	Bran	7.62 mg/100g	(Călinoiu & Vodnar, 2020)
Vitexin	Tartary Buckwheat (Kitawase, Hokkai T8,T9,T10)	Sprouts	45-645 mg/100g	(Arasu, <i>et al.</i> , 2014)
	Tartary Buckwheat	Flowers, leaves, stalk, root	0.33-4.22 mg/100g	(Dziedzic, <i>et al.</i> , 2018)
	Common Buckwheat	Hull	0.69-1.2 mg/100g	(Kalinova, <i>et al.</i> , 2019)
	Common Buckwheat/Tartary Buckwheat (Kitawase, Hokkai T9)	Seeds, Sprouts	20-630 mg/100g	(Kim, <i>et al.</i> , 2008)
	Common Buckwheat	Seeds	0.25-3.28 mg/100g	(Kiprovski, <i>et al.</i> , 2015)
	Tartary Buckwheat	flour	1-3 mg/100g	(Peng L., 2017)
	Tartary Buckwheat	bran	4-9 mg/100g	(Peng L., 2017)
	Common Buckwheat	Sprouts	463 mg/100g	(Chen, <i>et al.</i> , 2020)
	Common Buckwheat	Sprouts, microgreens, leafy greens	581.27 mg/100g	(Sharma, 2011)
isovitexin	Common Buckwheat	Sprouts, microgreens, leafy greens	370.14 mg/100g	(Sharma, 2011)
Avenanthramides				
Compound	Plant (Cultivar)	Plant parts	Concentration	Author
Total avenanthramides (AVAs)	Oat ( Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	4.20-9.10 mg/100g <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (13 cultivars)	Germ, endosperm, bran, hull	0.50-21.43 mg/100g	(Chen, <i>et al.</i> , 2018)
	Oat (8 cultivars)	Germ, endosperm, bran, hull	1.12-8.39 mg/100g	(Multari, <i>et al.</i> , 2018)
	Oat (commercial)	Bran	0.79-13.3 mg/100g	(Hu, <i>et al.</i> , 2019)
	Oat	naked	3.73- 71.85mg/100g	{Tong, 2014 #155}
AVA 2c	Oat ( Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	1.58-4.49 mg/100g <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Matilda, Sang, Freja)	Germ, endosperm, bran, hull	74-115 nmol/g	(Dimberg, <i>et al.</i> , 2005)
	Oat (Gem, Vista)	Leaves, spikelet, germ, endosperm, bran, hull	4.02-7.8 nmol/g	(D. M. Peterson & Dimberg, 2008)
	Oat (10 commercial brands)	Germ, endosperm, bran, hull	1.1-1.6 mg/100g	(Pridal, <i>et al.</i> , 2018)
	Oat (10 commercial brands)	Bran	0.9-2.9 mg/100g	(Pridal, <i>et al.</i> , 2018)
	Oat (10 commercial brands), rolled oats	Germ, endosperm, bran, hull	0.7-2.1 mg/100g	(Pridal, <i>et al.</i> , 2018)
	Oat (13 cultivars)	Germ, endosperm, bran, hull	0.16-7.00 mg/100g	(Chen, <i>et al.</i> , 2018)
	Oat (8 cultivars)	Germ, endosperm, bran, hull	0.35-3.92 mg/100g	(Multari, <i>et al.</i> , 2018)
	Oat (commercial)	Bran	0.41 mg/100g	(Călinoiu & Vodnar, 2020)

	Oat	naked	1.85 mg/100g	{Tong, 2014 #155}
Tocols				

	Oat (commercial), thermally processed	Bran	0.63 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (commercial)	Bran	0.07-1.49 mg/100g	(Hu, <i>et al.</i> , 2019)
	Oat (20 varieties)	Seeds	0.66 mg/100g	(Varga, <i>et al.</i> , 2018)
	Oat (20 varieties)	Hull	6.37 mg/100g	(Varga, <i>et al.</i> , 2018)
	Oat	Hulls, groats	0.11 mg/100g	(Ortiz-Robledo, <i>et al.</i> , 2013)
	Oat	Flakes, whole grain	0.4-0.9 mg/100g	(Mattila, <i>et al.</i> , 2005);
	Oat	grain	0.6-2.9 mg/100g	(Dokuyucu, <i>et al.</i> , 2003)
AVA 2p	Oat ( Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	1.3-3.0 mg/100g <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Matilda, Sang, Freja)	Germ, endosperm, bran, hull	78-120 nmol/g	(Dimberg, <i>et al.</i> , 2005)
	Oat (Gem, Vista)	Leaves, spikelet, germ, endosperm, bran, hull	3.93-74.26 nmol/g	(D. M. Peterson & Dimberg, 2008)
	Oat (10 commercial brands)	Germ, endosperm, bran, hull	0.6-1.5 mg/100g	(Pridal, <i>et al.</i> , 2018)
	Oat (10 commercial brands)	Bran	0.7-1.6 mg/100g	(Pridal, <i>et al.</i> , 2018)
	Oat (10 commercial brands), rolled oats	Germ, endosperm, bran, hull	0.6-1.9 mg/100g	(Pridal, <i>et al.</i> , 2018)
	Oat (13 cultivars)	Germ, endosperm, bran, hull	0.23-7.45 mg/100g	(Chen, <i>et al.</i> , 2018)
	Oat (8 cultivars)	Germ, endosperm, bran, hull	0.60-3.96 mg/100g	(Multari, <i>et al.</i> , 2018)
	Oat	Hulls, groats	0.15 mg/100g	(Ortiz-Robledo, <i>et al.</i> , 2013)
	Oat (commercial)	Bran	0.70 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (commercial), thermally processed	Bran	0.87 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (commercial)	Bran	0.06-3.32 mg/100g	(Hu, <i>et al.</i> , 2019)
	Oat	Flakes, whole grain	0.4-0.9 mg/100g	(Mattila, <i>et al.</i> , 2005);
	Oat	grain	0.3-1.9 mg/100g	(Dokuyucu, <i>et al.</i> , 2003)
	Oat (20 varieties)	Seeds	2.56 mg/100g	(Varga, <i>et al.</i> , 2018)
	Oat (20 varieties)	Hull	2.48 mg/100g	(Varga, <i>et al.</i> , 2018)
	Oat	Hulls, groats	0.12 mg/100g	(Ortiz-Robledo, <i>et al.</i> , 2013)
	Oat ( Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	1.27-2.40 mg/100g <sup>a</sup>	(Sterna, <i>et al.</i> , 2014)
	Oat (Matilda, Sang, Freja)	Germ, endosperm, bran, hull	62-65 nmol/g	(Dimberg, <i>et al.</i> , 2005)
	Oat (Gem, Vista)	Leaves, spikelet, germ, endosperm, bran, hull	3.35-18.16 nmol/g	(D. M. Peterson & Dimberg, 2008)
AVA 2f	Oat (10 commercial brands)	Germ, endosperm, bran, hull	0.9-2.3 mg/100g	(Pridal, <i>et al.</i> , 2018)
	Oat (10 commercial brands)	Bran	1.0-2.8 mg/100g	(Pridal, <i>et al.</i> , 2018)
	Oat (10 commercial brands), rolled oats	Germ, endosperm, bran, hull	0.6-2.2 mg/100g	(Pridal, <i>et al.</i> , 2018)
	Oat (13 cultivars)	Germ, endosperm, bran, hull	0.11-6.97 mg/100g	(Chen, <i>et al.</i> , 2018)
	Oat (8 cultivars)	Germ, endosperm, bran, hull	0.47-2.19 mg/100g	(Multari, <i>et al.</i> , 2018)
	Oat (commercial)	Bran	0.78 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (commercial), thermally processed	Bran	0.91 mg/100g	(Călinoiu & Vodnar, 2020)
	Oat (commercial)	Bran	0.05-6.62 mg/100g	(Hu, <i>et al.</i> , 2019)
	Oat	Flakes, whole grain	0.4-0.9 mg/100g	(Mattila, <i>et al.</i> , 2005)
	Oat	grain	0.3-2.8 mg/100g	(Dokuyucu, <i>et al.</i> , 2003)
	Oat (20 varieties)	Seeds	2.18 mg/100g	(Varga, <i>et al.</i> , 2018)
	Oat (20 varieties)	Hull	3.03 mg/100g	(Varga, <i>et al.</i> , 2018)

Compound	Plant	Plant part	Concentration	Author
Total tocols	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	1.61-3.61 mg/100g <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (12 cultivars) <sup>b</sup>	Germ, endosperm, bran	1.9 to 3.0 mg /100g <sup>c</sup>	(D.M. Peterson & Qureshi, 1993)
	Oat (unspecified) <sup>d</sup>	Germ, endospore, bran	7.21 mg/100g	(Panfili, <i>et al.</i> , 2003)
	Oat (Slawko)	Germ, endospore, bran	0.5 mg/100g	(Zielinski, <i>et al.</i> , 2001)
	Buckwheat Kora	whole grain	5.46 mg/100g	(Zielinski, <i>et al.</i> , 2001)
	Buckwheat Kora,	endosperm with embryo	6.44 mg/100g	(Zielinski, <i>et al.</i> , 2001)
	Buckwheat Mancan variety	seed	55.22mg/100g	(Przybylski, <i>et al.</i> , 1998)
Trienols	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	1.09-2.57 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat (Slawko)	Germ, endospore	1.35 mg/100g	(Zielinski, <i>et al.</i> , 2001)
$\alpha$ -T	Buckwheat	whole grain, hull, groat	3.32-9.44 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat Kora	whole grain	0.09 mg/100g	(Zielinski, <i>et al.</i> , 2001)
	Buckwheat Kora	endosperm with embryo	0.19 mg/100g	(Zielinski, <i>et al.</i> , 2001)
	Buckwheat Mancan variety	seed	38.01 mg/100g	(Przybylski, <i>et al.</i> , 1998)
	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	0.45-0.98 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat (12 cultivars) <sup>b</sup>	Germ, endosperm, bran	0.72 – 0.94 mg/100g <sup>c</sup>	(D.M. Peterson & Qureshi, 1993)
	Oat (unspecified) <sup>d</sup>	Germ, endospore, bran	1.49 mg/100g	(Panfili, <i>et al.</i> , 2003)
	Oat (Slawko)	Germ, endospore	0.09 mg/100g	(Zielinski, <i>et al.</i> , 2001)
	Oat (BDMY6, BDMY7, Che-chois, Y2330)	Germ, endosperm, bran, hull	0.56-4.14 mg/100g	(Musa Ozcan, <i>et al.</i> , 2006)
$\alpha$ -T3	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	0.94-2.30 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat (12 cultivars) <sup>b</sup>	Germ, endosperm, bran	0.09- 0.19 mg/100g <sup>c</sup>	(D.M. Peterson & Qureshi, 1993)
	Oat (unspecified) <sup>d</sup>	Germ, endospore, bran	5.64 mg/100g	(Panfili, <i>et al.</i> , 2003)
	Oat (Slawko)	Germ, endospore	0.27 mg/100g	(Zielinski, <i>et al.</i> , 2001)
$\beta$ -T	Buckwheat Mancan variety	seed	1.65 mg/100g	(Przybylski, <i>et al.</i> , 1998)
	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	0.06-0.10 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat (12 cultivars) <sup>b</sup>	Germ, endosperm, bran	0.07-0.13 mg/100g <sup>c</sup>	(D.M. Peterson & Qureshi, 1993)
	Oat (unspecified) <sup>d</sup>	Germ, endospore, bran	0.3 mg/100g	(Panfili, <i>et al.</i> , 2003)
	Oat (Slawko)	Germ, endospore	0.08 mg/100g	(Zielinski, <i>et al.</i> , 2001)
	Oat (BDMY6, BDMY7, Che-chois, Y2330)	Germ, endosperm, bran, hull	0.08-0.84 mg/100g	(Musa Ozcan, <i>et al.</i> , 2006)
$\beta$ -T3	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	0.13-0.27 mg/100g	(Shewry, <i>et al.</i> , 2008)
	Oat (12 cultivars) <sup>b</sup>	Germ, endosperm, bran	0.05- 0.15 mg/100g <sup>c</sup>	(D.M. Peterson & Qureshi, 1993)
	Oat (unspecified) <sup>d</sup>	Germ, endospore, bran	0.54 mg/ 100g	(Panfili, <i>et al.</i> , 2003)
	Oat (Slawko)	Germ, endospore	0.11 mg/100g	(Zielinski, <i>et al.</i> , 2001)
$\gamma$ -T	Buckwheat	whole grain, hull, groat	10.1-22.10 mg/100g	(Sedej, <i>et al.</i> , 2012)
	Buckwheat Kora	whole grain	5.14 mg/100g	(Zielinski, <i>et al.</i> , 2001)
	Buckwheat Kora	endosperm with embryo	6.04 mg/100g	(Zielinski, <i>et al.</i> , 2001)
	Buckwheat Mancan variety	seed	10.56 mg/100g	(Przybylski, <i>et al.</i> , 1998)
	Oat (12 cultivars) <sup>b</sup>	Germ, endosperm, bran	0.05-0.12 mg/100g <sup>c</sup>	(D.M. Peterson & Qureshi, 1993)
	Oat (unspecified) <sup>d</sup>	Germ, endospore, bran	0.04 mg/100g	(Panfili, <i>et al.</i> , 2003)
	Oat (BDMY6, BDMY7, Che-chois, Y2330)	Germ, endosperm, bran, hull	0.03-0.36 mg/100g	(Musa Ozcan, <i>et al.</i> , 2006)
$\delta$ - T	Buckwheat Kora	whole grain	0.24 mg/100g	(Zielinski, <i>et al.</i> , 2001)
	Buckwheat	endosperm with embryo	0.22 mg/100g	(Zielinski, <i>et al.</i> , 2001)

	Kora			
	Buckwheat Mancan variety	seed,	3.64 mg/100g	(Przybylski, <i>et al.</i> , 1998)
	Buckwheat	whole grain, hull, groat	0.37-0.46 mg/100g	(Sedej, <i>et al.</i> , 2012)
*Expressed originally as micrograms per gram, converted to milligram per 100g				

Supplemental table 3. Phytosterols, polysaccharides and fatty acids in oat and buckwheat				
Phytosterols				
Compound	Plant (cultivar)	Plant part	Concentration	Author
Total sterols	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	61.8-68.2 mg/100g <sup>a</sup>	(Horbowicz & Obendorf, 1992)
	Oat (Elin, Freja, Galopp, Matilda, Strok, Stormogul, Vital)	Germ, endosperm, bran, hull	35-49.1 mg/100g	{Maatta, 1999 #83}
	Common Buckwheat	Hulls, Groat	19-139 mg/100g	(Dziedzic, <i>et al.</i> , 2018)
	Buckwheat	Flour	99 mg/100g	{Bacchetti, 2011 #189}
Stanols, %	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	1.3-1.6 mg/100g <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
stanols	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	1.3-1.6 mg/100g <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
sitosterol	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	36.5-44.2mg/100g <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Vatter)	Leaves	42.6% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Vatter)	Germ, Endosperm, bran, hull	42.9% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Vatter)	Germ and endosperm	30.6% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Elin, Freja, Galopp, Matilda, Strok, Stormogul, Vital)	Germ, endosperm, bran, hull	23.7-32.1 mg/100g	{Maatta, 1999 #83}
campesterol	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	5.0-6.2 mg/100g <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Vatter)	Leaves	Trace	(Eichenberger, 1984)
	Oat (Vatter)	Germ, Endosperm, bran, hull	Trace	(Eichenberger, 1984)
	Oat (Vatter)	Germ, endosperm	trace	(Eichenberger, 1984)
	Oat (Vatter)	Leaves	Trace	(Eichenberger, 1984)
	Oat (Vatter)	Germ, Endosperm, bran, hull	Trace	(Eichenberger, 1984)
	Oat (Vatter)	Germ, endosperm	trace	(Eichenberger, 1984)
	Oat (Elin, Freja, Galopp, Matilda, Strok, Stormogul, Vital)	Germ, endosperm, bran, hull	3.2-4.6 mg/100g	{Maatta, 1999 #83}
stigmasterol	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	2.2-3.6 mg/100g <sup>a</sup>	(Peng L., 2017)
	Oat (Vatter)	Leaves	28.4% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Vatter)	Germ, Endosperm, bran, hull	3.9% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Vatter)	Germ, endosperm	4.6% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Elin, Freja, Galopp, Matilda, Strok, Stormogul, Vital)	Germ, endosperm, bran, hull	1.1-2.1 mg/100g	{Maatta, 1999 #83}
$\Delta^7$ - stigmasterol	Oat (Vatter)	Leaves	1.7% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Vatter)	Germ, Endosperm, bran, hull	1.0% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Vatter)	Germ, endosperm	2.0% <sup>b</sup>	(Eichenberger, 1984)
stigmasterol	Oat (Vatter)	Leaves	0.8%	(Eichenberger, 1984)
lophenol	Oat (Vatter)	Leaves	1.8%	(Eichenberger, 1984)
	Oat (Vatter)	Germ, Endosperm, bran, hull	---	(Eichenberger, 1984)
	Oat (Vatter)	Germ, endosperm	0.6%	(Eichenberger, 1984)
$\Delta^5$ -avenasterol	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	2.4- 3.4 mg/100g <sup>a</sup>	
	Oat (Vatter)	Leaves	2.4% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Vatter)	Germ, Endosperm, bran, hull	26.1% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Vatter)	Germ, endosperm	23.2% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Elin, Freja, Galopp, Matilda, Strok, Stormogul, Vital)	Germ, endosperm, bran, hull	1.5-4.7 mg/100g	{Maatta, 1999 #83}
$\Delta^7$ -avenastenol	Oat (Cacko, MV-Pehely, Fengli, Expander, Bajka)	Germ, endosperm, bran, hull (except for naked Cacko)	0.8- 1.2 mg/100g <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Vatter)	Leaves	1.7% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Vatter)	Germ, Endosperm, bran, hull	0.6% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Vatter)	Germ, endosperm	9.0% <sup>b</sup>	(Eichenberger, 1984)
cholestanol	Oat (Vatter)	Leaves	2.9% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Vatter)	Germ, Endosperm, bran, hull	Trace	(Eichenberger, 1984)
	Oat (Vatter)	Germ, endosperm	0.4% <sup>b</sup>	(Eichenberger, 1984)
$\Delta^7$ -cholestenol	Oat (Vatter)	Leaves	1.7% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Vatter)	Germ, Endosperm, bran, hull	0.6% <sup>b</sup>	(Eichenberger, 1984)
	Oat (Vatter)	Germ, endosperm	0.9% <sup>b</sup>	(Eichenberger, 1984)
<p>*Cacko is a naked oat variety with high protein content. It has a high soluble dietary fibre (including <math>\beta</math>-glucan) content, but it is also sensitive to the environment and may give lower yields than other cultivars. Bajka is an intensive type with good protein content and seed yield. Expander is also an intensive type with good yield and is less sensitive to environmental effects. MV-Pehely has high protein content, whereas Fengli is a tall type.</p> <p>**total phenolics as galic acid equivalents</p> <p>*** Total phenol (gallic acid equivalent) and flavonoid (quercetin equivalent) contents were determined using Folin–Ciocalteu and aluminum chloride reagents, respectively</p>				

**** catechin equivalents <sup>a</sup> Original units converted from microgram/gram to milligram/100g ((A.R. Gulpinar, <i>et al.</i> , 2011)) <sup>b</sup> Expressed in percent (%) of the total sterol extracted				
Polysaccharides				
Compound	Plant	Species/ cultivars	Concentration	Author
Total fagopyritols	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Whole Achenesa	128.2-962.1mg/100g	(Steadman, <i>et al.</i> , 2001)
fagopyritol A1	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Whole Achenesa	12.4-235.8 mg/100g	(Steadman, <i>et al.</i> , 2001)
fagopyritol A2	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Whole Achenesa	9.2-101.5mg/100g	(Steadman, <i>et al.</i> , 2001)
fagopyritol A3	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Whole Achenesa	11.1-65.8 mg/100g	(Steadman, <i>et al.</i> , 2001)
fagopyritol B1	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Whole Achenesa	78.9-1510.7 mg/100g	(Steadman, <i>et al.</i> , 2001)
fagopyritol B2	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Whole Achenesa	16.7-84.5 mg/100g	(Steadman, <i>et al.</i> , 2001)
fagopyritol B3	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Whole Achenesa	trace	(Steadman, <i>et al.</i> , 2001)
Fagopyritols	Buckwheat	bran	2600mg/100g	(Brindzova, <i>et al.</i> , 2008)
Total fagopyritols	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Groats	176.4-2774.5 mg/100g	(Steadman, <i>et al.</i> , 2001)
fagopyritol A1	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Groats	18.8-327.5mg/100g	(Steadman, <i>et al.</i> , 2001)
fagopyritol A2	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Groats	14.2-188.8mg/100g	(Steadman, <i>et al.</i> , 2001)
fagopyritol A3	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Groats	11.2-92.3mg/100g	(Steadman, <i>et al.</i> , 2001)
fagopyritol B1	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Groats	115.7mg/100g-2004 mg/100g	(Steadman, <i>et al.</i> , 2001)
fagopyritol B2	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Groats	16.5-216mg/100g	(Steadman, <i>et al.</i> , 2001)
fagopyritol B3	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Groats	trace	(Steadman, <i>et al.</i> , 2001)
D-chiro-inositol	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Groats	15.4-161.7mg/100g	(Steadman, <i>et al.</i> , 2001)
D-chiro-inositol	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Whole Achenesa	6.9-82.1mg/100g	(Steadman, <i>et al.</i> , 2001)
amylopectin	Buckwheat	Starch	75000 mg/100g	(Qin, <i>et al.</i> , 2010 ; Takahama & S., 2010)
myo-inositol	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Groats	8.4-88.1mh/100g	(Steadman, <i>et al.</i> , 2001)
myo-inositol	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Whole Achenesa	3.3-40.5 mg/100g	(Steadman, <i>et al.</i> , 2001)
galactinol	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Groats	5.2-70.4mg/g	(Steadman, <i>et al.</i> , 2001)
galactinol	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Whole Achenesa	3.5-36.5mg/100g	(Steadman, <i>et al.</i> , 2001)
sucrose	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Groats	392.6-2463.5mg/100g	(Steadman, <i>et al.</i> , 2001)
sucrose	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Whole Achenesa	223-3876.3mg/100g	(Steadman, <i>et al.</i> , 2001)
$\alpha$ -galactosides	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Groats	37-44%	(Steadman, <i>et al.</i> , 2001)
$\alpha$ -galactosides	Buckwheat	Buckwheat (Cv. Manor) Milling Fractions from Whole Achenesa	38-42%	(Steadman, <i>et al.</i> , 2001)
amylose	Buckwheat	starch	25000 mg/100g	(Qin, <i>et al.</i> , 2010; Takahama & S., 2010)
Phytic acid	Buckwheat	bran	3500-	(Sterna, <i>et al.</i> , 2014);

			3800mg/100 g	
Glucan	Oat (unspecified) <sup>b</sup>	Germ, endosperm, bran	6.3g/60g of starch	(Regand, <i>et al.</i> , 2011)
Beta glucan	Oat		3.9%	(Kourimska, <i>et al.</i> , 2018)
	Oat	flour	0.5-5.6%	(Van den Broeck, <i>et al.</i> , 2015)
	Oat (Cacko)	Germ, endosperm	1.09% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (MV-Pehely)	Germ, endosperm	1.12% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Fengli)	Germ, endosperm	1.03% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Expander)	Germ, endosperm	1.03% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Bajka)	Germ, endosperm	0.96% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Cacko)	Bran	8.33% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (MV-Pehely)	Bran	8.09% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Fengli)	Bran	6.20% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Expander)	Bran	8.18% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Bajka)	Bran	8.39% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Cacko)	Germ, endosperm, bran	5.5% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (MV-Pehely)	Germ, endosperm, bran	5.6% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Fengli)	Germ, endosperm, bran	4.5% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Expander)	Germ, endosperm, bran	5.1% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Bajka)	Germ, endosperm, bran	5.3% <sup>a</sup>	(Shewry, <i>et al.</i> , 2008)
	Oat (Cyril)	Germ, endosperm, bran	28 200 mg/100 g	(Hozova, <i>et al.</i> , 2007)
	Oat (Jakub)	Germ, endosperm, bran	5 180 mg/100 g	(Hozova, <i>et al.</i> , 2007)
	Oat (unspecified)	Groat	4.1-5.7%	(Sowa & White, 1992)
	Buckwheat Fagopyrum	Whole grain	1 260 mg/100g	(Hozova, <i>et al.</i> , 2007)
Fatty acids				
Total fatty acids	Buckwheat	bran	11000mg/100 g	(Steadman, <i>et al.</i> , 2001)
	Buckwheat	embryo	8.2% <sup>a</sup>	(Dorrell, 1971)
	Buckwheat	endosperm	0.4% <sup>a</sup>	(Dorrell, 1971)
	Buckwheat	testa	2.0% <sup>a</sup>	(Dorrell, 1971)
	Buckwheat	pericarp	0.5% <sup>a</sup>	(Dorrell, 1971)
	Common Buckwheat	Whole seed, hulls, bran, flour	288-5429mg/100g	(Sinkovic, <i>et al.</i> , 2020)
	Tartary Buckwheat	Whole seed, hulls, bran, flour	294-6264 mg/100g	(Sinkovic, <i>et al.</i> , 2020)
C14:0	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Flour	11-12.4 mg/100g	(Peng L., 2017)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Bran	23-23.4 mg/100g	(Peng L., 2017)
	Common Buckwheat	Embryo, Endosperm	0.3-14.9% <sup>a</sup>	(Horbowicz & Obendorf, 1992)
	Common Buckwheat	Whole seed, hulls, bran, flour	14-157 mg/100g	(Sinkovic, <i>et al.</i> , 2020)
	Tartary Buckwheat	Whole seed, hulls, bran, flour	14-81mg/100g	(Sinkovic, <i>et al.</i> , 2020)
	Common Buckwheat	grains	530-800mg/100g	(Golijan, <i>et al.</i> , 2019)
	Tartary Buckwheat	whole seed	0.1-0.5% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	embryo	10.8-17.4% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	endosperm	14.9-20% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	testa	10.4-18.2% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	pericarp	11% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Flour	184.2-238.9 mg/100g	(Peng L., 2017)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Bran	380.6-423.8 mg/100g	(Peng L., 2017)
	Common Buckwheat	Embryo, Endosperm	13.1-70.5% <sup>a</sup>	(Horbowicz & Obendorf, 1992)
	Common Buckwheat/Tartary Buckwheat	Seed	17.1-18.6% <sup>a</sup>	(Tsuzuki, <i>et al.</i> , 1991)
	Common Buckwheat	Whole seed, hulls, bran, flour	1473-2642 mg/100g	(Sinkovic, <i>et al.</i> , 2020)
	Tartary Buckwheat	Whole seed, hulls, bran, flour	1386-2434	(Sinkovic, <i>et al.</i> , 2020)

			mg/100g	
	Common Buckwheat	grains	18600-2401 mg/100g	(Golijan, <i>et al.</i> , 2019)
C18:0	Common Buckwheat	grains	4400-5090 mg/100g	(Golijan, <i>et al.</i> , 2019)
	Tartary Buckwheat	whole seed	2-2.4% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	embryo	1.6-2.2% <sup>a</sup>	(Dorrell, 1971)
	Tartary, Buckwheat	endosperm	2.2-3.7% <sup>a</sup>	(Dorrell, 1971)
	Tartary, Buckwheat	testa	2.2-2.8% <sup>a</sup>	(Dorrell, 1971)
	Tartary, Buckwheat	pericarp	2.5-5% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Flour	67.4-100.4 mg/100g	(Peng L., 2017)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Bran	105.4-125.4 mg/100g	(Peng L., 2017)
	Common Buckwheat	Embryo, Endosperm	2.6-5.3% <sup>a</sup>	(Horbowicz & Obendorf, 1992)
	Common Buckwheat/ Tartary Buckwheat	Seed	1.89-2.05% <sup>a</sup>	(Tsuzuki, <i>et al.</i> , 1991)
	Common Buckwheat	Whole seed, hulls, bran, flour	167-336 mg/100g	(Sinkovic, <i>et al.</i> , 2020)
	Tartary Buckwheat	Whole seed, hulls, bran, flour	193-499 mg/100g	(Sinkovic, <i>et al.</i> , 2020)
	Tartary Buckwheat	whole seed	32.5-38.5% <sup>a</sup>	(Dorrell, 1971)
	Common Buckwheat	grains	52020-53420 mg/100g	(Golijan, <i>et al.</i> , 2019)
	Tartary Buckwheat	embryo	33.2-40.5% <sup>a</sup>	(Dorrell, 1971)
	Tartary,Buckwheat	endosperm	31-35.5% <sup>a</sup>	(Dorrell, 1971)
	Tartary,Buckwheat	testa	28.7-33.9% <sup>a</sup>	(Dorrell, 1971)
	Tartary, Buckwheat	pericarp	14.6-19.6% <sup>a</sup>	(Dorrell, 1971)
	Common Buckwheat/Tartary Buckwheat	Seed	35.9-36.7% <sup>a</sup>	(Tsuzuki, <i>et al.</i> , 1991)
	Common Buckwheat	Embryo, Endosperm	6.1-33.7% <sup>a</sup>	(Horbowicz & Obendorf, 1992)
	Common Buckwheat	Whole seed, hulls, bran, flour	2096-3647mg/100g	(Sinkovic, <i>et al.</i> , 2020)
	Tartary Buckwheat	Whole seed, hulls, bran, flour	2620-4076 mg/100g	(Sinkovic, <i>et al.</i> , 2020)
C18:1	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Flour	264.7-283.9 mg/100g	(Peng L., 2017)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Bran	547.2-722.2 mg/100g	(Peng L., 2017)
	Common Buckwheat	grains	32740-53420 mg/100g	(Golijan, <i>et al.</i> , 2019)
C18:2	Common Buckwheat	grains	9330-12040mg/100 g	(Golijan, <i>et al.</i> , 2019)
	Common Buckwheat	Embryo,Endosperm	1.9-49.6% <sup>a</sup>	(Horbowicz & Obendorf, 1992)
	Common Buckwheat/Tartary Buckwheat	Seed	34.4-36.9% <sup>a</sup>	(Tsuzuki, <i>et al.</i> , 1991)
	Tartary, Buckwheat	whole seed	30.9-34.7% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	embryo	32.4-36.8% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	endosperm	16.8-33% <sup>a</sup>	(Dorrell, 1971)



	Tartary Buckwheat	, testa	27.6-32.7% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	, pericarp	13-25.7% <sup>a</sup>	(Dorrell, 1971)
	Common Buckwheat	Whole seed, hulls, bran, flour	3723-4757 mg/100g	(Sinkovic, <i>et al.</i> , 2020)
	Tartary Buckwheat	Whole seed, hulls, bran, flour	3554-4490mg/100g	(Sinkovic, <i>et al.</i> , 2020)
C18:2n-6c	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Flour	258.3-329.9 mg/100g	(Peng L., 2017)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Bran	793-1032.4 mg/100g	(Peng L., 2017)
C18:3n-3	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Flour	21.9-26.1mg/100g	(Peng L., 2017)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Bran	38.4-49.4mg/100g	(Peng L., 2017)
C18:3	Common Buckwheat	grains	5340-6740 mg/100g	(Golijan, <i>et al.</i> , 2019)
	Tartary Buckwheat	, whole seed	3.2-6-2% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	, embryo	2.4-6.5% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	, endosperm	2.6-5.7% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	, testa	4.4-6-4% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	, pericarp	2.9-4.2% <sup>a</sup>	(Dorrell, 1971)
	Common Buckwheat/Tartary Buckwheat	Seed	1.56-2.24% <sup>a</sup>	(Tsuzuki, <i>et al.</i> , 1991)
	Common Buckwheat	Embryo, Endosperm	1.1-12.9% <sup>a</sup>	(Horbowicz & Obendorf, 1992)
	Common Buckwheat	Whole seed, hulls, bran, flour	221-578 mg/100g	(Sinkovic, <i>et al.</i> , 2020)
	Tartary Buckwheat	Whole seed, hulls, bran, flour	168-360mg/100g	(Sinkovic, <i>et al.</i> , 2020)
C20:0	Common Buckwheat	grains	2720-3380 mg/100g	(Golijan, <i>et al.</i> , 2019)
	Tartary Buckwheat	whole seed	1.5-2.2% <sup>a</sup>	(Dorrell, 1971; A.R. Gulpinar, <i>et al.</i> , 2011)
	Tartary Buckwheat	embryo	1.1-1.6% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	endosperm	1.1-3-1% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	testa	1-3.3% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	pericarp	3.3-5-3% <sup>a</sup>	(Dorrell, 1971)
	Common Buckwheat/ Tartary Buckwheat	Seed	1.13-1.40% <sup>a</sup>	(Tsuzuki, <i>et al.</i> , 1991)
	Common Buckwheat	Embryo, Endosperm	1.1-2.4% <sup>a</sup>	(Horbowicz & Obendorf, 1992)
	Common Buckwheat	Whole seed, hulls, bran, flour	184-304 mg/100g	(Sinkovic, <i>et al.</i> , 2020)
	Tartary Buckwheat	Whole seed, hulls, bran, flour	113-203 mg/100g	(Sinkovic, <i>et al.</i> , 2020)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Flour	15.2-16.5 mg/100g	(Peng L., 2017)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Bran	34.4-35.6 mg/100g	(Peng L., 2017)
	Common Buckwheat	grains	0190-360 mg/100g	(Golijan, <i>et al.</i> , 2019)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Flour	8.8-12.1 mg/100g	(Peng L., 2017)
	Tartary Buckwheat Chuanqiao 1&2, Xiqiao 3	Bran	26.0-31.8 mg/100g	(Peng L., 2017)

	Common Buckwheat	Embryo, Endosperm	1.1-3.6% <sup>a</sup>	(Horbowicz & Obendorf, 1992)
	Common Buckwheat/ Tartary Buckwheat	Seed	2.12-2.95% <sup>a</sup>	(Tsuzuki, <i>et al.</i> , 1991)
C22:1	Tartary Buckwheat	Seed	0.500 mg/100g	Tsuzuki, <i>et al.</i> , 1991)
C22:0	Tartary Buckwheat	, whole seed	1.1-2.2% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	, embryo	0.8-1.7% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	, endosperm	0.9-2.4% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	, testa	1.2-4% <sup>a</sup>	(Dorrell, 1971)
	Tartary Buckwheat	, pericarp	0.1-6.2% <sup>a</sup>	(Dorrell, 1971)
	Common Buckwheat	Embryo, Endosperm	0.9-2.8% <sup>a</sup>	(Horbowicz & Obendorf, 1992)
	Common Buckwheat/Tartary Buckwheat	Seed	1.06-1.44% <sup>a</sup>	(Tsuzuki, <i>et al.</i> , 1991)
	Common Buckwheat	grains	250 mg/100g	(Golijan, <i>et al.</i> , 2019)
	Common Buckwheat/Tartary Buckwheat	Seed	0.18-0.52% <sup>a</sup>	(Tsuzuki, <i>et al.</i> , 1991)
	Common Buckwheat/Tartary Buckwheat	Seed	0.60-0.91% <sup>a</sup>	(Tsuzuki, <i>et al.</i> , 1991)
1,2 diacylglycerol	Oat (33 wild species)	Germ, endosperm, bran	1.5, % <sup>b</sup>	(Leonova, <i>et al.</i> , 2008)
	Oat (10 cultivated species)	Germ, endosperm, bran	1.1, % <sup>b</sup>	(Leonova, <i>et al.</i> , 2008)
1,3 diacylglycerol	Oat (33 wild species)	Germ, endosperm, bran	2.2, % <sup>b</sup>	(Leonova, <i>et al.</i> , 2008)
	Oat (10 cultivated species)	Germ, endosperm, bran	2.3, % <sup>b</sup>	(Leonova, <i>et al.</i> , 2008)
Free fatty acid	Oat (33 wild species)	Germ, endosperm, bran	1.6, % <sup>b</sup>	(Leonova, <i>et al.</i> , 2008)
	Oat (10 cultivated species)	Germ, endosperm, bran	2.4, % <sup>b</sup>	(Leonova, <i>et al.</i> , 2008)
	Oat	Groat, bran, endosperm, scutellum, embrioni	2-5%	(Youngs, 1978)
TAG1	Oat (wild species)	Germ, endosperm, bran	1.2, % <sup>b</sup>	(Leonova, <i>et al.</i> , 2008)
	Oat (cultivated species)	Germ, endosperm, bran	1.4, % <sup>b</sup>	(Leonova, <i>et al.</i> , 2008)
	Common buckwheat	grain	9.94-29.69 %	(Golijan, <i>et al.</i> , 2019)
TAG2	Oat (wild species)	Germ, endosperm, bran	0.9, % <sup>b</sup>	(Leonova, <i>et al.</i> , 2008)
	Oat (cultivated species)	Germ, endosperm, bran	1.1, % <sup>b</sup>	(Leonova, <i>et al.</i> , 2008)
Total Triacylglycerol	Oat (wild species)	Germ, endosperm, bran	77.1, % <sup>b</sup>	(Leonova, <i>et al.</i> , 2008)
	Oat (cultivated species)	Germ, endosperm, bran	74.2, % <sup>b</sup>	(Leonova, <i>et al.</i> , 2008)
	Oat (15 cultivars)	Groat, bran, endosperm, scutellum, embrioni	39-58%	(Youngs, 1978)
Myristic (C14:0)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	0.5-0.7%	(Kourimska, <i>et al.</i> , 2018)
	Oat (Laima, Arta, St.Liva, St.Darta)	Germ, endosperm, bran, hull	0.2%	(Sterna, <i>et al.</i> , 2014)
	Oat (15 cultivars)	Groat, bran, endosperm, scutellum, embrioni	0.4-0.8%	
	Oat (9 cultivars)	Germ, endosperm, bran, +/-hull	0.2-0.3%	(Brindzova, <i>et al.</i> , 2008)
Palmitic (C16:0)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, hull	21.4-22.8%	(Kourimska, <i>et al.</i> , 2018)
	Oat	flour	31.8- 262.3 mg/100g	(Van den Broeck, <i>et al.</i> , 2015)
	Common Buckwheat	Seeds	13.5%	(A. R. Gulpinar, <i>et al.</i> , 2012)

	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, hull	16.1-17.1%	(Sterna, <i>et al.</i> , 2014)
	Oat (S-156, 33793)	Germ, endosperm, bran	15.5-17.4%	(Sterna, <i>et al.</i> , 2014)
	Oat (15 cultivars)	Groat, bran, endosperm, scutellum, embrioni	16.1-21.8%	(Youngs, 1978)
	Oat (Astor, Maldwyn, C 5020, C5027, C5031, Nuprime)	Germ, endosperm, bran, +/-hull	16.6-21.7%	(Welch, 1975)
	Oat (9 cultivars)	Germ, endosperm, bran, +/-hull	13.8-17.2%	(Brindzova, <i>et al.</i> , 2008)
	Oat (Freja, Matilda)	Germ, endosperm, bran, hull	12.0-16.1%	(Banas, <i>et al.</i> , 2007)
	Oat (Freja, Matilda)	Emrbyo and scutellum	14.0-20.9%	(Banas, <i>et al.</i> , 2007)
	Oat (BDMY6, BDMY7, Che-chois, Y2330)	Germ, endosperm, bran, hull	15.7-18.8%	(Musa Ozcan, <i>et al.</i> , 2006)
Palmitoleic (C16:1 cis-9)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	0.2-0.3%	(Kourimska, <i>et al.</i> , 2018)
	Oat (9 cultivars)	Germ, endosperm, bran, +/-hull	0.2-0.3%	(Brindzova, <i>et al.</i> , 2008)
Stearic (C18:0)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	1.5-3-1%	(Kourimska, <i>et al.</i> , 2018)
	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, hull	1.7-2.5%	(Sterna, <i>et al.</i> , 2014)
	Oat	flour	7.6-33.3 mg/100g	(Van den Broeck, <i>et al.</i> , 2015)
	Common Buckwheat	Seeds	1.62%	(A. R. Gulpinar, <i>et al.</i> , 2012)
	Oat (S-156, 33793)	Germ, endosperm, bran	1.7-2.4%	(Sterna, <i>et al.</i> , 2014)
	Oat (15 cultivars)	Groat, bran, endosperm, scutellum, embrioni	1.2-2%	(Youngs, 1978)
	Oat (Astor, Maldwyn, C 5020, C5027, C5031, Nuprime)	Germ, endosperm, bran, +/-hull	1.05-2.18%	(Welch, 1975)
	Oat (9 cultivars)	Germ, endosperm, bran, +/-hull	1.3-2.2%	(Brindzova, <i>et al.</i> , 2008)
	Oat (Freja, Matilda)	Germ, endosperm, bran, hull	0.5-2.6%	(Banas, <i>et al.</i> , 2007)
	Oat (Freja, Matilda)	Emrbyo and scutellum	0.4-0.9%	(Banas, <i>et al.</i> , 2007)
	Oat (BDMY6, BDMY7, Che-chois, Y2330)	Germ, endosperm, bran, hull	2.79%	(Musa Ozcan, <i>et al.</i> , 2006)
Oleic (C18:1)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, hull	36.2-37.8%	(Sterna, <i>et al.</i> , 2014)
	Oat (S-156, 33793)	Germ, endosperm, bran	37.9-40.4%	(Sterna, <i>et al.</i> , 2014)
	Oat (Freja, Matilda)	Germ, endosperm, bran, hull	17.3-51.6%	(Banas, <i>et al.</i> , 2007)
	Oat (Freja, Matilda)	Emrbyo and scutellum	30.5-43.0%	(Banas, <i>et al.</i> , 2007)
	Oat (BDMY6, BDMY7, Che-chois, Y2330)	Germ, endosperm, bran, hull	33.97-51.26%	(Musa Ozcan, <i>et al.</i> , 2006)
	Common Buckwheat	Seeds	33.15%	(A. R. Gulpinar, <i>et al.</i> , 2012)
Elaidic (C18:1trans-9)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	0.1-0.5%	(Kourimska, <i>et al.</i> , 2018)
Oleic (C18:1 cis-9)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	30.7-32.2%	(Kourimska, <i>et al.</i> , 2018)
	Oat (15 cultivars)	Groat, bran, endosperm, scutellum, embrioni	28.4-40.3%	(Youngs, 1978)

	Oat (Astor, Maldwyn, C 5020, C5027, C5031, Nuprime)	Germ, endosperm, bran, +/-hull	22.0-32.5%	(Welch, 1975)
	Oat (9 cultivars)	Germ, endosperm, bran, +/-hull	33.4-42.2%	(Brindzova, <i>et al.</i> , 2008)
	Oat	flour	113.7-723.3 mg/100g	(Van den Broeck, <i>et al.</i> , 2015)
Cis-13-Octadecenoic (C18:1 cis-13)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	1.5-1.8%	(Kourimska, <i>et al.</i> , 2018)
Linoleic (C18:2)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, hull	1-1.2%	(Sterna, <i>et al.</i> , 2014)
	Oat	flour	1.2-17.1 mg/100g	(Van den Broeck, <i>et al.</i> , 2015)
	Oat (S-156, 33793)	Germ, endosperm, bran	0.9-1.2%	(Sterna, <i>et al.</i> , 2014)
	Oat (Astor, Maldwyn, C 5020, C5027, C5031, Nuprime)	Germ, endosperm, bran, +/-hull	44.4-50.3%	(Welch, 1975)
	Oat (9 cultivars)	Germ, endosperm, bran, +/-hull	34.4-42.7%	(Brindzova, <i>et al.</i> , 2008)
	Oat (Freja, Matilda)	Germ, endosperm, bran, hull	29.9-47.0%	(Banas, <i>et al.</i> , 2007)
	Oat (Freja, Matilda)	Emrbyo and scutellum	34.6-45.6%	(Banas, <i>et al.</i> , 2007)
	Oat (BDMY6, BDMY7, Che-chois, Y2330)	Germ, endosperm, bran, hull	22.8-35.9%	(Musa Ozcan, <i>et al.</i> , 2006)
	Common Buckwheat	Seeds	31.93%	(A. R. Gulpinar, <i>et al.</i> , 2012)
C18:2 (n-6)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, hull	39.4-41.6%	(Sterna, <i>et al.</i> , 2014)
	Oat (S-156, 33793)	Germ, endosperm, bran	38.4-39.1%	(Sterna, <i>et al.</i> , 2014)
Linoelaidic trans (C18:2 trans,trans-9,12)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	<0.1-0.4%	(Kourimska, <i>et al.</i> , 2018)
Linoelaidic cis (C18:2 cis,cis-9,12)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	34.6-38.2%	(Kourimska, <i>et al.</i> , 2018)
	Oat (15 cultivars)	Groat, bran, endosperm, scutellum, embrioni	36.6-45.8%	(Youngs, 1978)
Linolenic (C18:3)	Oat (15 cultivars)	Groat, bran, endosperm, scutellum, embrioni	1.5-2.5%	(Youngs, 1978)
	Oat (Astor, Maldwyn, C 5020, C5027, C5031, Nuprime)	Germ, endosperm, bran, +/-hull	1.89-3.38%	(Welch, 1975)
	Oat (9 cultivars)	Germ, endosperm, bran, +/-hull	1.1-1.9%	(Brindzova, <i>et al.</i> , 2008)
	Oat (Freja, Matilda)	Germ, endosperm, bran, hull	1.0-20.2%	(Banas, <i>et al.</i> , 2007)
	Oat (Freja, Matilda)	Emrbyo and scutellum	2.0-5.3%	(Banas, <i>et al.</i> , 2007)
	Oat (BDMY6, BDMY7, Che-chois, Y2330)	Germ, endosperm, bran, hull	0.64%	(Musa Ozcan, <i>et al.</i> , 2006)
Linolenic (C18:3 (n-3))	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, hull	0.9-1.3%	(Sterna, <i>et al.</i> , 2014)
	Oat (S-156, 33793)	Germ, endosperm, bran	0.9-1.1%	(Sterna, <i>et al.</i> , 2014)
Alpha-Linolenic (C18:3 all cis 9,12,15)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	1.2-1.6%	(Kourimska, <i>et al.</i> , 2018)
Linolenic (C18:3) isomers	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	0.1-0.4%	(Kourimska, <i>et al.</i> , 2018)

Arachidic (C20:0)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	0.2-0.4%	(Kourimska, <i>et al.</i> , 2018)
	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, hull	0.2%	(Sterna, <i>et al.</i> , 2014)
	Oat (S-156, 33793)	Germ, endosperm, bran	0.2%	(Sterna, <i>et al.</i> , 2014)
Paullinic (C20:1)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, hull	0.6-0.8%	(Sterna, <i>et al.</i> , 2014)
	Oat (S-156, 33793)	Germ, endosperm, bran	0.7%	(Sterna, <i>et al.</i> , 2014)
	Oat (9 cultivars)	Germ, endosperm, bran, +/-hull	0.8-1.0%	(Brindzova, <i>et al.</i> , 2008)
	Oat (Freja, Matilda)	Germ, endosperm, bran, hull	0.7-1.2%	(Banas, <i>et al.</i> , 2007)
	Oat (Freja, Matilda)	Emrbyo and scutellum	0.7-1.7%	(Banas, <i>et al.</i> , 2007)
Eicosapentanoic (C20:5)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, hull	0.2-0.4%	(Sterna, <i>et al.</i> , 2014)
	Oat (S-156, 33793)	Germ, endosperm, bran	0.2-0.3%	(Sterna, <i>et al.</i> , 2014)
Gondoic (C20:1 cis-11)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	0.8-1.1%	(Kourimska, <i>et al.</i> , 2018)
Eicosadienoic (C20:2 cis,cis-11,14)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	<0.1-1.5%	(Kourimska, <i>et al.</i> , 2018)
Behenic (C22:0)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	0.3-0.9%	(Kourimska, <i>et al.</i> , 2018)
	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, hull	0.1%	(Sterna, <i>et al.</i> , 2014)
	Oat (S-156, 33793)	Germ, endosperm, bran	0.1%	(Sterna, <i>et al.</i> , 2014)
Erucic (C22:1 cis-13)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	<0.1-0.1%	(Kourimska, <i>et al.</i> , 2018)
Lignoceric (C24:0)	Oat (Raven, Cavaliere, Korok, Kertag, Kamil, Patrik)	Germ, endosperm, bran, +/-hull	0.3-0.5%	(Hamberg, <i>et al.</i> , 1998)
Avenoleic acid	Oat (Vital)	Germ, endosperm, bran, hull (Eluent chloroform)	29% <sup>c</sup>	(Hamberg, <i>et al.</i> , 1998)
	Oat (Vital)	Germ, endosperm, bran, hull (Eluent Acetone)	63% <sup>c</sup>	(Hamberg, <i>et al.</i> , 1998)
	Oat (Vital)	Germ, endosperm, bran, hull (Eluent Methanol)	8% <sup>c</sup>	(Hamberg, <i>et al.</i> , 1998)
Galactolipid –avenolate complex	Oat (Vital)	Germ, endosperm, bran, hull	0.5-0.6mg/g	(Hamberg, <i>et al.</i> , 1998)
Cholesterol	Oat (Vatter)	Leaves	12.8%	(Eichenberger, 1984)
	Oat (Vatter)	Germ, Endosperm, bran, hull	2.8%	(Eichenberger, 1984)
	Oat (Vatter)	Germ and endosperm	2.4%	(Eichenberger, 1984)
Saponins				
Avenacoside A	oat	grains	0.15-2.6 mg/100g	(Pecio, <i>et al.</i> , 2013)
	oat	kernel	0.008-0.03% DW	(Önning & Asp, 1993)
avenacoside B	oat	grains	0.06-2.51 mg/100g	(Pecio, <i>et al.</i> , 2013)
	oat	kernel	0.003-0.01% DW	(Önning & Asp, 1993)
26-desglucoavenacoside A	oat	grains	0.07-0.7 mg/100g	(Pecio, <i>et al.</i> , 2013)

Total avenacins	oat	roots	1280 mg/100g	(Mary & Crombie, 1986)
Total saponins	oat	Porridge	130 mg/100g	(Tschesche & Wulff, 1973)
	oat	Porridge	100 mg/100g	(Fenwick D. E, 1983)

## References

- Alvarez-Jubete, L., Wijngaard, H., Arendt, E. K., & Gallagher, E. (2010). Polyphenol composition and in vitro antioxidant activity of amaranth, quinoa buckwheat and wheat as affected by sprouting and baking. *Food Chemistry*, 119(2), 770-778.
- Andrea Brunori, G. S., Hao Xi, Gerardo Baviello, Bernadett Nehiba, Gyula Rabnecz, György Végvári. (2009). Rutin Content of the Grain of 22 Buckwheat (*Fagopyrum esculentum* Moench and *Fagopyrum tataricum* Gaertn.) Varieties Grown in Hungary *The European Journal of Plant Science and Biotechnology*.
- Arasu, M. V., Kim, S. J., Al-Dhabi, N. A., Suzuki, T., Yamauchi, H., & Lee, S. W. (2014). Comparison of Flavonoid Contents between Common and Tartary Buckwheat (*Fagopyrum*) Sprouts Cultured with/without Soil. *Asian Journal of Chemistry*, 26(18), 5985-5990.
- Bai, C. Z., Feng, M. L., Hao, X. L., Zhong, Q. M., Tong, L. G., & Wang, Z. H. (2015). Rutin, quercetin, and free amino acid analysis in buckwheat (*Fagopyrum*) seeds from different locations. *Genetics and Molecular Research*, 14(4), 19040-19048.
- Banas, A., Debski, H., Banas, W., Heneen, W. K., Dahlqvist, A., Bafor, M., Gummesson, P. O., Marttila, S., Ekman, A., Carlsson, A. S., & Stymne, S. (2007). Lipids in grain tissues of oat (*Avena sativa*): differences in content, time of deposition, and fatty acid composition. *J Exp Bot*, 58(10), 2463-2470.
- Brindzova, L., Certik, M., Raptá, P., Zalibera, M., Mikulajova, A., & Takacsova, M. (2008). Antioxidant activity, beta-glucan and lipid contents of oat varieties. *Czech Journal of Food Sciences*, 26(3), 163-173.
- Călinoiu, L. F., & Vodnar, D. C. (2020). Thermal processing for the release of phenolic compounds from wheat and oat bran. *Biomolecules*, 10(1).
- Chen, C., Li, T., Chen, Z., Wang, L., & Luo, X. (2020). Absorption Rates and Mechanisms of Avenanthramides in a Caco-2 Cell Model and Their Antioxidant Activity during Absorption. *J Agric Food Chem*, 68(8), 2347-2356.
- Chen, C., Wang, L., Wang, R., Luo, X., Li, Y., Li, J., Li, Y., & Chen, Z. (2018). Phenolic contents, cellular antioxidant activity and antiproliferative capacity of different varieties of oats. *Food Chem*, 239, 260-267.
- Danila, A. M., Kotani, A., Hakamata, H., & Kusu, F. (2007). Determination of rutin, catechin, epicatechin, and epicatechin gallate in buckwheat *Fagopyrum esculentum* Moench by micro-high-performance liquid chromatography with electrochemical detection. *J Agric Food Chem*, 55(4), 1139-1143.
- Dimberg, L. H., Gissen, C., & Nilsson, J. (2005). Phenolic compounds in oat grains (*Avena sativa* L.) grown in conventional and organic systems. *Ambio*, 34(4-5), 331-337.
- Dokuyucu, T., Peterson, D., & Akkaya, A. (2003). Contents of Antioxidant Compounds in Turkish Oats: Simple Phenolics and Avenanthramide Concentrations. *Cereal Chemistry - CEREAL CHEM*, 80, 542-543.
- Dorrell, D. G. (1971). Fatty Acid Composition of Buckwheat Seed. *Journal of the American Oil Chemists Society*, 48(11), 693-8.
- Dziadek, K., Kopeć, A., Pastucha, E., Piątkowska, E., Leszczyńska, T., Pisulewska, E., Witkiewicz, R., & Francik, R. (2016). Basic chemical composition and bioactive compounds content in selected cultivars of buckwheat whole seeds, dehulled seeds and hulls. *Journal of Cereal Science*, 69.
- Dziedzic, K., Drozdzyńska, A., Gorecka, D., & Czaczyk, K. (2009). Contents of some selected antioxidants in buckwheat and products produced during its processing. *Zywnosc-Nauka Technologia Jakosc*, 16(6), 81-90.
- Dziedzic, K., Gorecka, D., Szwengiel, A., Sulewska, H., Kreft, I., Gujska, E., & Walkowiak, J. (2018). The Content of Dietary Fibre and Polyphenols in Morphological Parts of Buckwheat (*Fagopyrum tataricum*). *Plant Foods for Human Nutrition*, 73(1), 82-88.
- Eichenberger, W. B. (1984). Sterols in seeds and leaves of oats (*Avena sativa* L.). *Plant Cell Reports. Plant Cell Reports*, 3(6), 226-229.
- Fabjan, N., Rode, J., Kosir, I. J., Wang, Z., Zhang, Z., & Kreft, I. (2003). Tartary buckwheat (*Fagopyrum tataricum* Gaertn.) as a source of dietary rutin and quercitrin. *J Agric Food Chem*, 51(22), 6452-6455.
- Fenwick D. E, O. D. (1983). Saponin content of food plants and some prepared foods. *Journal of the Science of Food and Agriculture*, 34(2), 186-191.
- Gallagher, R. S., Ananth, R., Granger, K., Bradley, B., Anderson, J. V., & Fuerst, E. P. (2010). Phenolic and short-chained aliphatic organic acid constituents of wild oat (*Avena fatua* L.) seeds. *J Agric Food Chem*, 58(1), 218-225.
- Garleb, K. A., Bourquin, L. D., Hsu, J. T., Wagner, G. W., Schmidt, S. J., & Fahey, G. C., Jr. (1991). Isolation and chemical analyses of nonfermented fiber fractions of oat hulls and cottonseed hulls. *J Anim Sci*, 69(3), 1255-1271.

- Ge, R. H., & Wang, H. (2020). Nutrient components and bioactive compounds in tartary buckwheat bran and flour as affected by thermal processing. *International Journal of Food Properties*, 23(1), 127-137.
- Golijan, J., Milinčić, D., Petronijevic, R., Pešić, M., Barac, M., Sečanski, M., Iekić, S., & Kostić, A. (2019). The fatty acid and triacylglycerol profiles of conventionally and organically produced grains of maize, spelt and buckwheat. *Journal of Cereal Science*, 90, 102845.
- Gulpinar, A. R., Orhan, I. E., Kan, A., Senol, F. S., Celik, S. A., & Kartal, M. (2011). Estimation of in vitro neuroprotective properties and quantification of rutin and fatty acids in buckwheat (*Fagopyrum esculentum* Moench) cultivated in Turkey. *Food Research International*, 46(2), 536-543.
- Gulpinar, A. R., Orhan, I. E., Kan, A., Senol, F. S., Celik, S. A., & Kartal, M. (2012). Estimation of in vitro neuroprotective properties and quantification of rutin and fatty acids in buckwheat (*Fagopyrum esculentum* Moench) cultivated in Turkey. *Food Research International*, 46(2), 536-543.
- Habtemariam, S. (2019). Antioxidant and Rutin Content Analysis of Leaves of the Common Buckwheat (*Fagopyrum esculentum* Moench) Grown in the United Kingdom: A Case Study. *Antioxidants*, 8(6), 10.
- Hamberg, M., Liepinsh, E., Otting, G., & Griffiths, W. (1998). Isolation and structure of a new galactolipid from oat seeds. *Lipids*, 33(4), 355-363.
- Holasovaa, M., Fiedlerovaa, V., Smrcinovaa, H., Orsakb, M., Lachmanb, J., & Vavreinovaa, S. (2002). Buckwheat-the source of antioxidant activity in functional foods. *Food Research International*, 35(2-3), 207-211.
- Horbowicz, M., & Obendorf, R. L. (1992). Changes in Sterols and Fatty-Acids of Buckwheat Endosperm and Embryo during Seed Development. *J Agric Food Chem*, 40(5), 745-750.
- Hozova, B., Kuniak, L., Moravcikova, P., & Gajdošová, A. (2007). Determination of water-insoluble  $\beta$ -D-Glucan in the whole-grain cereals and pseudocereals. *Czech Journal of Food Sciences*, 25, 316-324.
- Hu, C., Tang, Y., Zhao, Y., & Sang, S. (2019). Quantitative Analysis and Anti-inflammatory Activity Evaluation of the A-Type Avenanthramides in Commercial Sprouted Oat Products. *J Agric Food Chem*.
- Hung, P. V., & Morita, N. (2008). Distribution of phenolic compounds in the graded flours milled from whole buckwheat grains and their antioxidant capacities. *Food Chem*, 109(2), 325-331.
- Jiang, P., Burczynski, F., Campbell, C., Pierce, G., Austria, J. A., & Briggs, C. J. (2007). Rutin and flavonoid contents in three Nutritional profile of buckwheat 17 buckwheat species *Fagopyrum esculentum*, *F. tataricum*, and *F. homotropicum* and their protective effects against lipid peroxidation. *Food Research International*(40), 356-364.
- Kalinova, J. P., Vrchotova, N., & Triska, J. (2019). Phenolics levels in different parts of common buckwheat (*Fagopyrum esculentum*) achenes. *Journal of Cereal Science*, 85, 243-248.
- Keriene, I., Mankeviciene, A., Bliznikas, S., Jablonskyte-Rasce, D., Maiksteniene, S., & Cesnuleviciene, R. (2015). Biologically active phenolic compounds in buckwheat, oats and winter spelt wheat. *Zemdirbyste-Agriculture*, 102(3), 289-296.
- Kim, S. J., Zaidul, I. S., Suzuki, T., Mukasa, Y., Hashimoto, N., Takigawa, S., Noda, T., Matsuura-Endo, C., & Yamauchi, H. (2008). Comparison of phenolic compositions between common and tartary buckwheat (*Fagopyrum*) sprouts. *Food Chem*, 110(4), 814-820.
- Kiprovski, B., Mikulic-Petkovsek, M., Slatnar, A., Veberic, R., Stampar, F., Malencic, D., & Latkovic, D. (2015). Comparison of phenolic profiles and antioxidant properties of European *Fagopyrum esculentum* cultivars. *Food Chemistry*, 185, 41-47.
- Kočevar Glavač, N., Stojilkovski, K., Kreft, S., Park, C., & Kreft, I. (2017). Determination of fagopyrins, rutin, and quercetin in Tartary buckwheat products. *LWT - Food Science and Technology*, 79.
- Kourimska, L., Sabolova, M., Horcicka, P., Rys, S., & Bozik, M. (2018). Lipid content, fatty acid profile, and nutritional value of new oat cultivars. *Journal of Cereal Science*, 84, 44-48.
- Kovacova, M., & Maliňová, E. (2007). Ferulic and coumaric acids, total phenolic compounds and their correlation in selected oat genotypes. *Czech Journal of Food Sciences*, 25(6), 325.
- Kraujaliene, V., Pukalskas, A., & Venskutonis, P. R. (2017). Multi-stage recovery of phytochemicals from buckwheat (*Fagopyrum esculentum* Moench) flowers by supercritical fluid and pressurized liquid extraction methods. *Industrial Crops and Products*, 107, 271-280.
- Kreft, S., Janeš, D., & Kreft, I. (2013). The content of fagopyrin and polyphenols in common and tartary buckwheat sprouts. *Acta Pharmaceutica*, 63(4), 553-560.
- Lee, J. M., Lee, K. H., Yoon, Y. H., Cho, E. J., & Lee, S. (2013). Identification of triterpenoids and flavonoids from the seeds of tartary buckwheat. *Natural Product Sciences*, 19(2), 137-144.
- Leonova, S., Shelenga, T., Hamberg, M., Konarev, A. V., Loskutov, I., & Carlsson, A. S. (2008). Analysis of oil composition in cultivars and wild species of oat (*Avena* sp.). *J Agric Food Chem*, 56(17), 7983-7991.
- Li, F. H., Yuan, Y., Yang, X. L., Tao, S. Y., & Ming, J. (2013). Phenolic Profiles and Antioxidant Activity of Buckwheat (*Fagopyrum esculentum* Moench and *Fagopyrum tartaricum* L. Gaerth) Hulls, Brans and Flours. *Journal of Integrative Agriculture*, 12, 1684-1693.



- Liu, Y., Cai, C., Yao, Y., & Xu, B. (2019). Alteration of phenolic profiles and antioxidant capacities of common buckwheat and tartary buckwheat produced in China upon thermal processing. *Journal of the Science of Food and Agriculture*, 99(12), 5565-5576.
- Mary, W., & Crombie, L. (1986). Distribution of avenacins A-1, A-2, B-1 and B-2 in oat roots: Their fungicidal activity towards 'take-all' fungus. *Phytochemistry*, 25, 2069-2073.
- Mattila, P., Pihlava, J.-M., & Hellström, J. (2005). Contents of Phenolic Acids, Alkyl- and Alkenylresorcinols, and Avenanthramides in Commercial Grain Products. *J Agric Food Chem*, 53, 8290-8295.
- Mazza, G. (1988). Lipid content and fatty acid composition of buckwheat seed. *Cereal Chemistry*, 65, 122-126.
- Multari, S., Pihlava, J. M., Ollennu-Chuasam, P., Hietaniemi, V., Yang, B., & Suomela, J. P. (2018). Identification and Quantification of Avenanthramides and Free and Bound Phenolic Acids in Eight Cultivars of Husked Oat (*Avena sativa* L) from Finland. *J Agric Food Chem*, 66(11), 2900-2908.
- Musa Ozcan, M., Ozkan, G., & Topal, A. (2006). Characteristics of grains and oils of four different oats (*Avena sativa* L.) cultivars growing in Turkey. *Int J Food Sci Nutr*, 57(5-6), 345-352.
- Önning, G., & Asp, N.-G. (1993). Analysis of saponins in oat kernels. *Food Chemistry*, 48, 301-305.
- Oomah, B. D., & Mazza, G. (1996). Flavonoids and Antioxidative Activities in Buckwheat. *J Agric Food Chem*, 44(7), 1746-1750.
- Ortiz-Robledo, F., Villanueva, I., Oomah, B. D., Lares-Asseff, I., Proal-Nájera, J., & Návar, J. (2013). Avenanthramides and nutritional components of four Mexican oat (*Avena sativa* L.) varieties. *Agrociencia*, 47, 225-232.
- Panfili, G., Fratianni, A., & Irano, M. (2003). Normal phase high-performance liquid chromatography method for the determination of tocopherols and tocotrienols in cereals. *J Agric Food Chem*, 51(14), 3940-3944.
- Park, C. H., Yeo, H. J., Park, Y. E., Chun, S. W., Chung, Y. S., Lee, S. Y., & Park, S. U. (2019). Influence of Chitosan, Salicylic Acid and Jasmonic Acid on Phenylpropanoid Accumulation in Germinated Buckwheat (*Fagopyrum esculentum* Moench). *Foods*, 8(5), 10.
- Pecio, Ł., Wawrzyniak-Szołkowska, A., Oleszek, W., & Stochmal, A. (2013). Rapid analysis of avenacosides in grain and husks of oats by UPLC-TQ-MS. *Food Chemistry*, 141, 2300-2304.
- Peng L., Z. L., Tan M., Deng Y., Yan J., Yan Z., Zhao G. (2017). Free amino acids, fatty acids and phenolic compounds in Tartary buckwheat of different hull colour. . *Czech Journal of Food Sciences*, 35, 214-222.
- Peterson, D. M., & Dimberg, L. H. (2008). Avenanthramide concentrations and hydroxycinnamoyl-CoA : hydroxyanthranilate N-hydroxycinnamoyltransferase activities in developing oats. *Journal of Cereal Science*, 47(1), 101-108.
- Peterson, D. M., & Qureshi, A. A. (1993). Genotype and environment effects on tocols of barley and oats. *Cereal Chemistry*(70), 157-162.
- Pomeranz, Y., & Robbins, G. S. (1972). Amino-Acid Composition of Buckwheat. *J Agric Food Chem*, 20(2), 270-&.
- Pridal, A. A., Bottger, W., & Ross, A. B. (2018). Analysis of avenanthramides in oat products and estimation of avenanthramide intake in humans. *Food Chem*, 253, 93-100.
- Przybylski, R., Lee, Y. C., & Eskin, N. A. M. (1998). Antioxidant and radical-scavenging activities of buckwheat seed components. *Journal of the American Oil Chemists Society*, 75(11), 1595-1601.
- Qin, P. Y., Wang, Q., Shan, F., Hou, Z. H., & Ren, G. X. (2010). Nutritional composition and flavonoids content of flour from different buckwheat cultivars. *International Journal of Food Science and Technology*, 45(5), 951-958.
- Quettier-Deleu, C., Gressier, B., Vasseur, J., Dine, T., Brunet, C., Luyckx, M., Cazin, M., Cazin, J. C., Bailleul, F., & Trotin, F. (2000). Phenolic compounds and antioxidant activities of buckwheat (*Fagopyrum esculentum* Moench) hulls and flour. *J Ethnopharmacol*, 72(1-2), 35-42.
- Regand, A., Chowdhury, Z., Tosh, S. M., Wolever, T. M. S., & Wood, P. (2011). The molecular weight, solubility and viscosity of oat beta-glucan affect human glycemic response by modifying starch digestibility. *Food Chem*, 129(2), 297-304.
- Rocchetti, G., Miragoli, F., Zacconi, C., Lucini, L., & Rebecchi, A. (2019). Impact of cooking and fermentation by lactic acid bacteria on phenolic profile of quinoa and buckwheat seeds. *Food research international (Ottawa, Ont.)*, 119, 886-894.
- Sedej, I., Sakac, M., Mandic, A., Misan, A., Tumbas, V., & Canadanovic-Brunet, J. (2012). Buckwheat (*Fagopyrum esculentum* Moench) grain and fractions: antioxidant compounds and activities. *Journal of Food Science*, 77(9), C954-959.
- Sharma, P. (2011). Phenolic contents, antioxidant and  $\alpha$ -glucosidase inhibition properties of Nepalese strain buckwheat vegetables. *African J Biotech*, 11.
- Shewry, P. R., Piironen, V., Lampi, A. M., Nystrom, L., Li, L., Rakszegi, M., Fras, A., Boros, D., Gebruers, K., Courtin, C. M., Delcour, J. A., Andersson, A. A., Dimberg, L., Bedo, Z., & Ward, J. L. (2008). Phytochemical and

- fiber components in oat varieties in the HEALTHGRAIN Diversity Screen. *J Agric Food Chem*, 56(21), 9777-9784.
- Sinkovic, L., Kokalj, D., Vidrih, R., & Meglic, V. (2020). Milling fractions fatty acid composition of common (Fagopyrum esculentum Moench) and tartary (Fagopyrum tataricum (L.) Gaertn) buckwheat. *Journal of Stored Products Research*, 85, 5.
- Siwatch, M., Yadav, R. B., & Yadav, B. S. (2019). Influence of processing treatments on nutritional and physicochemical characteristics of buckwheat (Fagopyrum esculentum). *Current Nutrition and Food Science*, 15(4), 408-414.
- Skoglund, M., Peterson, D., Andersson, R., Nilsson, J., & Dimberg, L. (2008). Avenanthramide content and related enzyme activities in oats as affected by steeping and germination. *Journal of Cereal Science*, 48, 294-303.
- Sowa, S. M. H., & White, P. J. (1992). Characterization of Starch Isolated from Oat Groats with Different Amounts of Lipid. *Cereal Chemistry*, 69(5), 521-527.
- Soycan, G., Schär, M. Y., Kristek, A., Boberska, J., Alsharif, S. N. S., Corona, G., Shewry, P. R., & Spencer, J. P. E. (2019). Composition and content of phenolic acids and avenanthramides in commercial oat products: Are oats an important polyphenol source for consumers? *Food Chemistry: X*, 3.
- Steadman, K. J., Burgoon, M. S., Lewis, B. A., Edwardson, S. E., & Obendorf, R. L. (2001). Minerals, phytic acid, tannin and rutin in buckwheat seed milling fractions. *Journal of the Science of Food and Agriculture*, 81(11), 1094-1100.
- Sterna, V., Zute, S., Brunava, L., & Vicupe, Z. (2014). Lipid Composition of Oat Grain Grown in Latvia. *9th Baltic Conference on Food Science and Technology - Food for Consumer Well-Being: Foodbalt 2014*, 77-80.
- Sytar, O. (2014). Phenolic acids in the inflorescences of different varieties of buckwheat and their antioxidant activity. *Journal of King Saud University - Science*, 27.
- Takahama, U., & S., H. (2010). Fatty acids, epicatechindimethylgallate, and rutin interact with buckwheat starch inhibiting its digestion by amylase: implications for the decrease in glycemic index by buckwheat flour. . *Agricultural and Food Chemistry*(58), 12431-12439.
- Terpinc, P., Cigic, B., Polak, T., Hribar, J., & Pozrl, T. (2016). LC-MS analysis of phenolic compounds and antioxidant activity of buckwheat at different stages of malting. *Food Chem*, 210, 9-17.
- Tien, N. N. T., Ngoc, D., Inoue, N., Morita, N., & Hung, P. (2018). Nutritional Composition, bioactive compounds and diabetic enzyme inhibition capacity of three varieties of buckwheat in Japan. *Cereal Chemistry*, 95.
- Tschesche, R., & Wulff, G. (1973). Chemie und Biologie der Saponine. In, vol. 30 (pp. 461-606).
- Tsuzuki, W., Ogata, Y., Akasaka, K., Shibata, S., & Suzuki, T. (1991). Fatty-acid composition of selected buckwheat species by fluorometric high-performance liquidchromatography. *Cereal Chemistry*, 68, 365-369.
- Van den Broeck, H., Londono, D., Timmer, R., Smulders, M. J. M., Gilissen, L. J. W. J., & Meer, I. (2015). Profiling of Nutritional and Health-Related Compounds in Oat Varieties. *Foods*, 5, 2.
- Varga, M., Jójárt, R., Fónad, P., Mihály, R., & Palágyi, A. (2018). Phenolic composition and antioxidant activity of colored oats. *Food Chemistry*, 268, 153-161.
- Wefers, D., & Bunzel, M. (2015). Characterization of Dietary Fiber Polysaccharides from Dehulled Common Buckwheat ( Fagopyrum esculentum) Seeds. *Cereal Chemistry*, 92(6), 598-603.
- Welch, R. W. (1975). Fatty acid composition of grain from winter and spring sown oats, barley and wheat. *Journal of the Science of Food and Agriculture*, 26(4), 429-435.
- Xing, Y. M., & White, P. J. (1997). Identification and function of antioxidants from oat greats and hulls. *Journal of the American Oil Chemists Society*, 74(3), 303-307.
- Youngs, V. L. (1978). Oat Lipids, *Cereal Chem.*, 55, 591-597.
- Zielińska, D., Turemko, M., Kwiatkowski, J., & Zieliński, H. (2012). Evaluation of flavonoid contents and antioxidant capacity of the aerial parts of common and tartary buckwheat plants. *Molecules (Basel, Switzerland)*, 17(8), 9668-9682.
- Zielinski, H., Ciska, E., & Kozłowska, H. (2001). The cereal grains: focus on vitamin E. *Czech Journal of Food Sciences*(19), 182-188.



# PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
<b>TITLE</b>			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
<b>ABSTRACT</b>			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2
<b>INTRODUCTION</b>			
Rationale	3	Describe the rationale for the review in the context of what is already known.	5
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5
<b>METHODS</b>			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	6
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	6
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	6
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Supplement
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	6
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	7
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	7
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	n.a.
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	n.a
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., $I^2$ for each meta-analysis).	n.a



# PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	n.a
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	n.a
<b>RESULTS</b>			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	7
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	7
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	n.a
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	n.a.
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	n.a.
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	n.a.
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	n.a.
<b>DISCUSSION</b>			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	7-17
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	17
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	18
<b>FUNDING</b>			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	19

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: [www.prisma-statement.org](http://www.prisma-statement.org).

**Declaration of interests**

☐ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☒ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

This research was supported by Standard Process. Hua Kern, Weston Bussler and Brandon Metzger are scientists at Standard Process Nutrition Innovation Center. Other authors have nothing to disclose.

## \*Credit Author Statement

\* Credit Author Statement

Study concept and design: Taulant Muka, Marija Glisic and Hua Kern.

Search strategy creation and online database search: Beatrice Minder.

Acquisition, collection, interpretation of data: Peter Francis Raguindin, Oche Adam Itodo, Jivko Stoyanov, Gordana M. Dejanovic, Magda Gamba, Eralda Asllanaj.

Drafting of the manuscript: Peter Francis Raguindin, Oche Adam Itodo, Marija Glisic and Hua Kern.

Critical revision of the manuscript for important intellectual content: Jivko Stoyanov, Gordana M. Dejanovic, Magda Gamba, Eralda Asllanaj, Beatrice Minder, Weston Bussler, Brandon Metzger, Taulant Muka.

Study supervision: Marija Glisic and Hua Kern.

All authors approved the final version of the manuscript.

Bern, 28-06-2018

Marija Glisic

A handwritten signature in dark blue ink, reading "M. Glisic". The signature is written in a cursive, flowing style. The letters are connected, with the "M" and "G" being particularly prominent. The signature is set against a light gray, textured background that resembles a piece of paper or a card.